

CUCURBITS (EXCEPT CA)

General Information

DuPont CURZATE 60DF must be used only in accordance with directions on the label.

PRODUCT INFORMATION

CURZATE 60DF is a locally systemic fungicide labeled in all states for the control of late blight on potatoes, potato seed pieces, and tomatoes and for downy mildew control on cucurbit crops, hop, lettuce (head and leafy), and spinach.

Potato seed pieces, tomatoes, cucurbits crops, hops: Except CA

CURZATE 60DF rapidly penetrates into plant tissues and is rainfast within 2 hours after application.

DO NOT use CURZATE 60DF alone. Always tank-mix with a labeled rate of protectant fungicide. Follow the more restrictive labeling of any tank mix partners. Do not tank mix with products that contain a prohibition on tank mixing.

Apply as a spray with ground, air, or chemigation (potatoes, cucurbits, lettuce, spinach, and tomatoes only) equipment, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Use only in commercial or farm plantings. Not for use in residential plantings or once any commercial crop is turned into U-Pick, Pick Your Own or similar operation.

RESISTANCE MANAGEMENT

Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. Therefore, tankmixing Curzate with a labeled rate of protectant fungicide that has a different mode of action is required. The tank mix partner must be labeled for downy mildew or late blight control. This ensures optimum performance and further reduces the potential for resistance development. For guidance on the particular crop and disease control situation, consult your state extension specialist or official state recommendations.

APPLICATION INFORMATION

Application Volumes

- For conventional ground application, apply a minimum of 20 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of foliage.
- For air-assisted ground application, apply a minimum of 10 gallons per acre.
- For aerial application, apply a minimum of 5 gallons per acre.

Chemigation

- For chemigation information for potatoes, cucurbits, lettuce, spinach, hops and tomatoes, see chemigation section following the crop-specific application directions.

Potato Seed Piece Treatment

- For potato seed piece treatment, see the Potato Seed Treatment section of the label.

Limitations, Restrictions, and Exceptions

CUCURBITS

Use CURZATE 60DF at 3.2 to 5 oz. per acre for control of downy mildew (*Pseudoperonospora cubensis*). Use only in combination with a labeled rate of a protectant fungicide, such as products containing mancozeb (e.g., DuPont MANZATE), maneb (e.g., DuPont MANEX), copper hydroxide (e.g., DuPont KOCIDE), or chlorothalonil.

Application Information

- Apply CURZATE 60DF plus a protectant fungicide on a 5-7 day schedule.
- Apply no more than 30 oz per 12 month period.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[N. A.](#)