

SOYBEAN - AERIAL WEB BLIGHT, ETC.

General Information

GENERAL INFORMATION

Bumper 41.8 EC is a broad spectrum fungicide for the control of specified diseases in labeled crops. Do not use this product in greenhouses or as a tree injection.

When an adjuvant is to be used with this product, Makhteshim Agan of North America, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

APPLICATION INSTRUCTIONS

For best results, sufficient water volume should be used to provide thorough coverage. In most situations, Bumper 41.8 EC is most effective when applied and allowed to dry before a rainfall. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

AERIAL APPLICATION: For those crops other than tree crops where aerial applications are indicated, apply in a minimum of 2 gals. of water per acre, unless specified otherwise in the APPLICATION INSTRUCTIONS section of the label. For tree crops, a minimum of 5 to 10 gals. Of water per acre is recommended using the higher volume on large trees unless specified otherwise in the APPLICATION INSTRUCTIONS

section of the label.

GROUND APPLICATION: For tree crops, a minimum of 50 gals. of water per acre is recommended unless specified otherwise in the APPLICATION INSTRUCTIONS section of the label. For all other crops, apply Bumper 41.8 EC by ground equipment in a minimum of 10 gals. of water per acre unless specified otherwise in the APPLICATION INSTRUCTIONS section of the label.

BANDED APPLICATION: For banded applications, the treated area is the area covered by the band, not total cropland planted.

RESISTANCE MANAGEMENT

Bumper 41.8 EC is effective in controlling pests and minimizing the development of resistance when used in rotation with other fungicides in an IPM program. To reduce selection pressure for resistant pests:

- Bumper 41.8 EC should be used in rotation with classes of fungicides with different modes of action.
- Use Bumper 41.8 EC as part of a pest management program that includes cultural and biological control where possible.
- Bumper 41.8 EC is in the Group 3 class of fungicides. The mode of action for propiconazole, the active ingredient in this product, is as a demethylation inhibitor of sterol biosynthesis (DMI) which disrupts membrane synthesis by blocking demethylation. Resistance can develop when products with the same mode of action are used repeatedly.
- Consult your State or local agricultural pest control advisor(s) for pest control strategies established for your area.

Refer to the label for tank mix information.

Limitations, Restrictions, and Exceptions

SOYBEAN

Applications may be made using ground or aerial application equipment. Use dilution rates found in the APPLICATION INSTRUCTIONS section of the label. When applying by air, adding an oil-based additive is recommended for improved coverage and penetration.

Apply 5 – 6 fl. oz. at the first appearance of Aerial web blight and repeat the application 14 to 21 days later. Under severe conditions, use the higher rate and shorter interval.

For control of other foliar diseases, apply 6 fl. oz. at growth stage R3 (early pod set)

when pods are 1/8 to 1/4 inch long and 21 days later at growth stage R5 (pod fill).

Apply 4 – 6 fl. oz. at first indication that soybean rust is in the area. For best control, preventative applications work best.

Repeat on a 14 to 21 day interval using the higher rate and shorter interval when disease is present in field and incidence is less than 2% (2 plants in 100 infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for rust and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications. On certain varieties, Bumper applications may cause crinkled or smaller greener leaves. Yields of dry beans displaying these characteristics have not been reduced due to propiconazole treatments.

- Applications may be made up to growth stage R6.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[At growth stage R3 \(early pod set\) when pods are 1/8 to 1/4 inch long and 21 days later at growth stage R5 \(pod fill\).](#)