

SOYBEAN - CUTWORM, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

SOYBEAN

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.

Apply with either aerial or ground equipment using sufficient spray volume to obtain full coverage of the plant and foliage. Use a minimum of 2 gallons of finished spray per acre by air or 10 gallons of finished spray per acre by ground. The addition of one to two quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control.

Do not make applications less than 7 days apart.

Do not graze or harvest treated soybean forage, straw, or hay for livestock feed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insect populations reaching locally determined economic threshold levels.](#)