

BENTGRASS TURF

General Information

GENERAL INFORMATION:

Trimec Encore Broadleaf Herbicide provides selective broadleaf control in warm season and cool season turfgrass. Established turf of warm season grasses include, but are not limited to, bermudagrass, bahiagrass, and zoysiagrass. Established turf of cool season grasses include, but are not limited to, Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra or established turf containing desirable clovers.

USE PRECAUTIONS - OBSERVE CAREFULLY.

1. Do not pour spray solutions near these plants.
2. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present.
3. Use only lawn type sprayers.
4. Avoid application to exposed active feeder roots of shallow-rooted trees such as poplar species and sensitive ornamentals. Do not exceed specified dosages for any area.
5. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
6. Do not spray bermudagrass or zoysiagrass while emerging from dormancy, unless injury can be tolerated.
7. Applications to newly seeded lawns may cause injury. Delay application until well established or after two or three mowings.
8. Do not broadcast apply when air temperatures exceed 90°F; some injury may be expected with spot treatments when air temperatures exceed 90°F.
9. Seed can be sown 3 to 4 weeks after application.
10. Do not contaminate domestic or irrigation waters.
11. Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass; only spot spray weeds on these types of grasses. Clover will be damaged if sprayed.
12. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.

13. Failure to observe the above precautions may result in injury to desirable plants.

14. Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

BENTGRASS TURF:

In closely-mowed bentgrass (putting and bowling greens) apply at a maximum rate of 1 fluid ounce of product per 1,500 square feet, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing bentgrass or injury may result. High spray volumes (i.e., 1 fluid ounce of product in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-operated sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy local overapplication and subsequent turf discoloration or injury.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[In May or mid-August through September.](#)