

PINEAPPLE - FLOWER INDUCTION

General Information

This Product:

Encourages faster coloration and maturity of APPLES, GRAPES, PEPPERS, and TOMATOES.

Loosens APPLES, CHERRIES, and WALNUTS for an earlier and more efficient harvest.

Encourages earlier, uniform coloring of mature FLUE-CURED TOBACCO.

Encourages fruit abscission (slipping) in CANTALOUPEs.

Augments hardiness in dormant fruit buds and helps to delay the spring bloom of SWEET CHERRIES in the Pacific Northwest.

Accelerates the loosening and ripening of BLACKBERRIES.

Minimizes lodging in WHEAT and BARLEY.

Eliminates leafy mistletoe from ORNAMENTAL DECIDUOUS TREES and dwarf mistletoe from ORNAMENTAL CONIFERS.

Removes unwanted fruit on APPLE, CRABAPPLE, CAROB, and OLIVE trees.

GENERAL INFORMATION

Contact your Extension Pomologist, Farm Advisor, Horticultural Specialist or Arysta LifeScience Representative for local recommendations on product spray volume, spray equipment and rates of application for varying weather conditions.

APPLICATION VOLUMES AND SPRAY COVERAGE

For optimum product efficacy, thorough spray coverage is necessary. This can be influenced by type of spray equipment, spray boom setup, nozzle selection, plant size, canopy density and spray pressure. Depending on these choices, the necessary spray volume will vary. For applications by air in California and Arizona, more than 5 gallons per acre must be used.

USE PRECAUTIONS

THE MIXTURE OF THIS PRODUCT WITH AMMONIUM THIOSULFATE IS PROHIBITED AS IT MAY CREATE TOXIC FUMES. Other than recommended on this label, this product should not be used with additives.

Upon mixture, this product should be applied as soon as possible; in no case should the spray solution be stored overnight.

Detrimental changes to plant growth, reduced yields, and plant injury may result from spray drift of this product to nearby crops and thus should be avoided. Do not plant another crop within 30 days after treatment.

This product is corrosive. Therefore, spills of concentrated product on the aircraft or other spray equipment should be avoided. Should such contact be made, immediately rinse with water.

EQUIPMENT CLEANING

This product is corrosive. As a result, spray deposit exposure will, over a period of time, damage metal, some paints and acrylic plastics. No more than one hour after exposure to spray deposits, these materials should be carefully rinsed with water and detergent.

Limitations, Restrictions, and Exceptions

PINEAPPLE

PINEAPPLE FLOWER INDUCTION: Application of ETHEPHON 2 will induce uniform flower initiation of pineapple plants.

- ETHEPHON 2 applied to pineapple plants will stimulate uniform initiation of flowering.

Timing of Application

- "Bearing age" about 12 months after planting. Generally about 6 months prior to desired harvest

Instructions

- Proper rate will vary with local growing conditions, varieties, plantation management practices and time of year. Use the higher rate where earlier harvest is desired. Apply when pineapple foliage is dry.

Comments

- Do not graze pineapple forage treated with ETHEPHON 2. Do not harvest

pineapples treated with ETHEPHON 2 sooner than 2 days after the last application.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

48 hours

The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

Timings

[\"Bearing age\" about 12 months after planting. Generally about 6 months prior to desired harvest.](#)