

# **NON-CROP AREAS - ANNUAL GRASS SUPPRESSION - FALL APPLICATION - AR, LA, MS AND TN**

## General Information

### GENERAL INFORMATION

Select 2 EC is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

Select 2 EC is a selective postemergence herbicide for control of annual and perennial grasses. Select 2 EC does not control sedges or broadleaf weeds.

Repeated use of Select 2 EC (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or applications conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies such as crop rotation, mechanical removal, and other classes of herbicides are not used from year to year.

Do not allow Select 2 EC to come in contact with desirable grass crops such as corn, rice, sorghum, small grains, or turf, as these and other grass crops will be injured or killed.

Minor leaf spotting may occur on treated plants under certain environmental conditions. New foliage is not affected.

### Control Symptoms

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days after application, depending on grass species treated and environmental conditions.

### APPLICATION INFORMATION

## Timing of Applications

Apply Select 2 EC postemergence to actively growing grasses according to rate table recommendations in the label.

Applications made to grass plants stressed by insufficient moisture, hot or cold temperatures, or to grass plants exceeding recommended growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, Select 2 EC should be applied as soon as possible after irrigation (within 7 days). In arid regions, a second application of Select 2 EC will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of Select 2 EC may reduce weed control. DO NOT APPLY Select 2 EC if rainfall is expected within one hour, since control may be reduced.

## Ground Application

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 40 gals. of spray solution per acre. Under the following conditions a minimum of 10 gals. per acre is required: ultra narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals. per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to onions (dry bulbs), garlic, and shallots (dry bulbs) should be made in a minimum of 20 gals. of spray solution per acre.

## Air Application

Use a minimum of 3 gals. of spray solution per acre unless otherwise directed in the label. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense. For onions (dry bulbs), garlic, or shallots (dry bulbs): When applying by air do

not exceed 8 fl. oz./A in a single application. In California, air applications to onions, garlic or shallots should be made in a minimum of 20 gals. of spray solution per acre. In states other than California, air application to onions, garlic, or shallots should be made in a minimum of 10 gals. of spray solution.

NOTE: Crop injury may occur when Select 2 EC is applied to onions, garlic, or shallots with aerial equipment.

### Spot Treatment

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/4% to 1/2% (0.33 oz. to 0.65 oz. per gal.) Select 2 EC and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz. per gal.) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz. per gal.) by volume.

NOTE: If Select 2 EC is applied as a spot treatment, care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.

## RESTRICTIONS AND LIMITATIONS

### GENERAL

Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory.

Do not apply a postemergence broadleaf herbicide within one day following application of Select 2 EC or reduced grass control may result.

Select 2 EC is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

For canola, clover, flax, mustard seed and radish crops, do not apply more than 16 fl. oz. of Select 2 EC (0.25 lb. ai) per acre per season. For all other crops, do not apply more than 32 fl. oz. of Select 2 EC (0.50 lb. ai) per season. Application on Long Island, New York, is restricted to no more than 16 fl. oz. of Select 2 EC (0.25 lb. ai) per acre per season.

Do not apply more than 8 fl. oz./A of Select 2 EC per application to the following crops: garden beets, carrots, radish, (and other root vegetables), leaf lettuce,

broccoli, cabbage, cauliflower (and other head & stem brassica vegetables), celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, flax and strawberry. Do not apply more than 6 fl. oz./A of Select 2 EC per application to canola, or mustard seed. Exceeding these recommendations may result in unacceptable crop injury.

Do not apply under conditions of stress. Applying Select 2 EC under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate Select 2 EC effectively, and will be less susceptible to herbicide activity.

Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, result in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than 2 Select 2 EC applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions.

Grass crops such as corn, rice, sorghum, small grains, or turf, etc. are highly sensitive to Select 2 EC.

While all the vegetable crops on the label have been tested and are tolerant to Select 2 EC, not all specialty varieties of these crops have been tested. It is advised that, before applying Select 2 EC to specialty varieties of vegetable crops on the label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.

Tank mixes of Select 2 EC and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of Select 2 EC may be necessary.

For tank mix application instructions, refer to the label.

## Limitations, Restrictions, and Exceptions

### NON-CROP AREAS

Annual Grasses that exceed heights claimed for control on the label. May be applied to grasses in the seed head stage (suppression only).

Important Target Weeds Include: Rhizome Johnsongrass, Bermudagrass, Tall Fescue and Ryegrass.

Adjuvant: SELECT 2 EC Herbicide must be applied with a crop oil concentrate at 1.0 qt./A.

General Information: Grass suppression only is being claimed on the label. A single application is recommended during the fall period. If regrowth occurs, the following spring a second application should be made. If at the time of the second application weed size is within label claims, then use the recommended label rate (6 to 8 fl. oz./A for annual grasses for control. Tillage after application may improve overall weed control. Do not mow or cultivate within 14 days of the application.

### Use Restrictions for SELECT 2 EC Herbicide:

1. Always read and follow all label directions when using any pesticide.
2. Do not plant any crop for 30 days after the application unless clethodim is registered for use in that crop.
3. Do not apply more than 32 fl. oz. of SELECT 2 EC Herbicide per acre per season.
4. Do not apply SELECT 2 EC Herbicide when grasses are under stress. Do not apply SELECT 2 EC if grassy weeds have been damaged by frost. Allow grass to resume new growth before application.

If grass has been mowed or cut as a function of the harvesting process, then grass should be allowed to resume new active growth before application.

5. Do not apply SELECT 2 EC Herbicide through any type of irrigation system.
6. Do not apply SELECT 2 EC Herbicide if rain is expected within 1 hour of

application.

7. Do not tank mix SELECT 2 EC Herbicide with other pesticides when using for grass suppression.

#### APPLICATION INFORMATION

For best results, SELECT 2 EC Herbicide should be applied to actively growing weeds.

Ground Application: Use of sufficient spray volume and pressure is essential to ensure complete coverage. Use a minimum of 10 gallons of spray solution per acre. Spray pressure should reflect a minimum of 40 psi and a maximum of 60 psi at the nozzle. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles for postemergence spray applications.

Spot Spray Application: When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/2% (0.65 oz. per gal.) SELECT 2 EC Herbicide and treat to wet vegetation, while not allowing runoff of spray solution. Add crop oil concentrate at 1% (1.3 oz. per gal.) by volume.

This 2(ee) bulletin contains directions for use which do not appear on the package label of SELECT 2 EC Herbicide. Follow instructions carefully.

Follow all applicable directions, restrictions and precautions on the EPA-registered labels.

#### RECOMMENDATIONS FOR ANNUAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

#### NOTE:

Weed Height: Generally occurs between 3-leaf stage and tillering.

Crabgrass, Crowfootgrass and Goosegrass: Length of lateral growth.

(1) Sethoxydim resistant volunteer corn.

(2) Includes Roundup Ready, Liberty Link and IMI-Corn volunteer corn.

(3) When the cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum Select 2 EC use rate for control is 8 fl. oz./A. Select 2 EC Herbicide.

(4) Rates higher than 8 fl. oz./A may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A may be applied. Do not apply more than 8 fl. oz./A of Select 2 EC per application to the following crops: garden beets, carrots, radish (and other root vegetables), leaf lettuce, broccoli, cabbage, cauliflower (and other head and stem brassica vegetables), celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, flax and strawberry. Do not apply more than 6 fl. oz./A of Select 2 EC per application to canola, or mustard seed.

For specific weed height in inches with its corresponding rates, refer to the label.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

#### Rates

[field\\_rates 0](#)

- 

#### Restricted Entry Interval

24 hours

Timings

Post-harvest

Postemergence (Weed)

Fall.