

ORNAMENTAL PLANTS - FLOWERING PLANTS AND BULBS - CARNATION, ETC. - LEAFSPOTS/FOLIAR BLIGHTS AND FLOWER SPOTS/BLIGHTS

General Information

PRODUCT USE INFORMATION

PRODUCT USE PRECAUTIONS AND RESTRICTIONS

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches and theme parks.

Do not apply to forests.

Agricultural Use Sites Only (sod farms, farms, forests, nurseries and greenhouses): This product must not be applied within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

DO NOT combine Daconil Zn in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. DO NOT combine Daconil Zn with Dipel, Latron B-1956 or Latron AG-98, horticultural oil, and products containing xylene as phytotoxicity may result from the combination when applied to some species on the label.

Limitations, Restrictions, and Exceptions

ORNAMENTAL PLANTS

Apply Daconil Zn at a rate of 2 pints (1.0 lb. a.i.) per 100 gallons of water unless other directions are given. DO NOT apply more than 70 pints Daconil Zn (36.4 lbs. a.i./acre) per growing season to field grown ornamentals. Apply in a spray to run-off, when conditions are favorable for disease development. Repeat applications at 7 to

14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply Daconil Zn at 7 day intervals. The minimum re-treatment interval is 7 days. Daconil Zn should be applied to plants when both foliage and flowers are dry, or nearly dry.

DO NOT combine Daconil Zn in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use.

Do not apply Daconil Zn through high pressure spray equipment. Do not use mistblowers, cold fog, or other fogging application equipment when making applications of Daconil Zn in greenhouses.

Do not apply Daconil Zn to either green or variegated Pittosporum or to Schefflera, as multiple applications may cause phytotoxic responses.

Do not apply Daconil Zn to ferns.

Ornamentals listed have been tested and found to tolerate applications of Daconil Zn at the recommended rates. The user should test for possible phytotoxic responses, using recommended rates on ornamental plants on a small area prior to commercial use. Applications made during bloom may damage flowers and/or fruits.

Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN.**

Flowering Plants: Avoid applications during bloom period on plants where flower injury is unacceptable.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

when both foliage and flowers are dry, or nearly dry.

When conditions are favorable for disease development.