

# **TURFGRASS - SNOW MOLD**

## General Information

### GENERAL USE PRECAUTIONS

HAWAII: Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures.

NASSAU AND SUFFOLK COUNTIES, NEW YORK: Use is limited to ornamentals grown in interiorscapes, greenhouses, lath and shade houses, containers, or other enclosed structures.

### RESISTANCE MANAGEMENT

Medallion is a protective fungicide used to aid in the control of soil-borne diseases. Medallion contains fludioxonil which is in the phenylpyrrole class of chemistry and has a unique mode of action, which prevents fungal respiration (Fungicide Action Group 12). Fungal isolates with acquired resistance to Group 12 may eventually dominate the fungal population if Group 12 fungicides are used repeatedly or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by fludioxonil or other Group 12 fungicides. A disease management program that includes alternation or tank mixes between Medallion and other labeled fungicides that have a different mode of action may prevent pathogen populations from developing resistance. Sanitation and other cultural practices to minimize disease are also recommended to aid in control, as well as to assist in preventing/delaying resistance development.

### APPLICATION INSTRUCTIONS

For applications of Medallion as soil drench applications only. Do not use in overhead irrigation systems.

### Limitations, Restrictions, and Exceptions

### TURFGRASS

Application Interval: Late Fall

Apply one to two applications in late fall before snow cover. Do not apply on top of

snow. For best disease control, tank mix with one to two registered snow mold products. For example, mix with a combination of Banner MAXX and Daconil Ultrex. Medallion at 0.25-0.3 oz. rate can be used when used when tank mixing with Banner MAXX and Daconil Ultrex when disease conditions are expected to be light to moderate.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

- 

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[In late fall before snow cover.](#)