

## **COLE CROPS - CABBAGE LOOPER, ETC.**

### General Information

#### GENERAL INFORMATION

ACEPHATE 75 SP is an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in ACEPHATE 75 SP is acephate, a water soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 75 SP through ingestion than by contact. Application of ACEPHATE 75 SP to maintain control should be repeated only as directed.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.**

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

#### Limitations, Restrictions, and Exceptions

#### ADDITIONAL USE INSTRUCTIONS

Apply in water at 25 to 150 gals. spray per acre by ground or a minimum of 5 gals. spray per acre by air.

Use the higher rate when heavy infestations of aphids are present.

Repeat application as needed but do not exceed maximum application rate of 2 2/3 lbs. of ACEPHATE 75 SP per acre per crop cycle.

Diamondback Moth Larvae — This insect has demonstrated an ability to develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current recommended control practices for this insect.

USE PRECAUTIONS:

- Do not feed trimmings to livestock or allow animals to graze in treated areas.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

24 hours

Timings

[When eggs or insects first appear.](#)