

PEANUTS - GRASSHOPPERS

General Information

GENERAL INFORMATION

ACEPHATE 75 SP is an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in ACEPHATE 75 SP is acephate, a water soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 75 SP through ingestion than by contact. Application of ACEPHATE 75 SP to maintain control should be repeated only as directed.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

Limitations, Restrictions, and Exceptions

ADDITIONAL USE INSTRUCTIONS

Apply in water at 10 to 50 gals. spray per acre by ground or in 5 to 10 gals. spray per acre by air.

Begin applications when eggs or insects first appear and repeat as needed to maintain control but do not exceed maximum application rate of 5 1/3 lbs. of ACEPHATE 75 SP per acre per crop cycle. ACEPHATE 75 SP can be tank mixed with at-cracking and early post emergence peanut herbicides, provided those products do not prohibit tank mixes, provided the most restrictive of label limitations and precautions are observed, and provided no label dosage rates are exceeded.

To determine physical compatibility, pour the recommended proportions of each chemical with the same proportion of water as will be present in the chemical

supply tank into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is considered physically compatible. When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides. Combinations should be kept agitated and should be applied immediately. Do not allow combinations to set for prolonged periods in the chemical supply tank.

USE PRECAUTIONS

- Do not feed treated forage or hay to livestock or allow animals to graze treated areas.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[When eggs or insects first appear.](#)