

ORNAMENTAL TREES AND SHRUBS - SCALE CRAWLERS

General Information

GENERAL INFORMATION

ACEPHATE 75 SP is an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in ACEPHATE 75 SP is acephate, a water soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 75 SP through ingestion than by contact. Application of ACEPHATE 75 SP to maintain control should be repeated only as directed.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

Limitations, Restrictions, and Exceptions

TIME OF APPLICATION

Repeat applications, at a 2 week or more interval, may be necessary where there is continuous crawler production.

RECOMMENDED APPLICATION

Apply the specified amount of ACEPHATE 75 SP in 100 gals. water with a hydraulic sprayer.

The addition of a suitable sticker improves control of Gypsy Moth larvae.

Mist blower application: Adjust rates to 1 1/3 lbs. per 100 gals. water for Gypsy Moth control and 1.0 lb. per 100 gals. water for Tent Caterpillar control.

- Do not apply to Huckleberry, Balm of Gilead, Cottonwood, Lombardy Poplar and Viburnum suspensum.

- Nursery crops. Before treating large plantings, spray only a few plants and observe two weeks for phytotoxicity.

Method

[Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[As crawlers begin to appear.](#)