

# **GREENHOUSE FLORAL AND FOLIAGE PLANT CROPS - SCALE CRAWLERS**

## General Information

### GENERAL INFORMATION

ACEPHATE 75 SP is an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in ACEPHATE 75 SP is acephate, a water soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 75 SP through ingestion than by contact. Application of ACEPHATE 75 SP to maintain control should be repeated only as directed.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

### Limitations, Restrictions, and Exceptions

#### TIME OF APPLICATION

Repeat applications, at a 2 week or more interval, may be necessary where there is continuous crawler production.

#### RECOMMENDED APPLICATION

Apply the specified amount of ACEPHATE 75 SP in 100 gals. water with a hydraulic sprayer as a full coverage spray. The addition of a wetting agent may be required on difficult to wet foliage.

Caution: Phytotoxicity has occurred on the following foliage plants: Bletchum gibbum, Cissus antarctica, Ficus triangularis, Fittonia verschaffeltii, Maranta

leuconeura kerchoveana, Plectranthus lutes, Plectranthus australis, Polipodium aureus, Polystichum. Pteris ensiformis, Tolmiea menziesii. Before treating large plantings spray only a few plants and observe 2 weeks for varietal phytotoxicity.

#### USE PRECAUTIONS:

- Do not apply more often than once every 28 days to Carnations and Chrysanthemum.
- CAUTION: Phytotoxicity has occurred on the following Chrysanthemum varieties: Albatross, Bonnie Jean, Dixie, Garland, Gent, Iceberg, Pride, Showoff, Statesman, Tally Ho, Westward Ho and Wild Honey. Before treating large Chrysanthemum plantings, spray only a few plants and observe two weeks for varietal phytotoxicity.
- Do not apply to chrysanthemums and roses with open flowers.
- Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

#### Method

[Spray](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

•

#### Restricted Entry Interval

24 hours

#### Timings

[As crawlers begin to appear.](#)