

# **FOR CONTROL OF GRASSES AND WEEDS IN NON-CROP AND INDUSTRIAL AREAS**

General Information

## **DIRECTIONS FOR USE**

### Application Restrictions:

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow others to enter the treated area until dusts have settled.

Total application of this product must not exceed 600 pounds of product per acre per year (12 pounds of bromacil per acre per year).

Application with a spoon, a pump-feed backpack spreader or a gravity-feed backpack spreader is prohibited.

Apply a maximum of 12 pounds active ingredient per acre in areas of high rainfall or dense vegetation.

Apply a maximum of 8 pounds active ingredient per acre in all other areas.

Apply a maximum of two applications per year. Minimum treatment interval: 90 days.

Not to be used in any recreational area or around homes.

Aerial application is prohibited.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. Al., v. EPA, C01-0132C (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/litstatus/wtc/index.htm>.

Limitations, Restrictions, and Exceptions

## NON-CROP WEED CONTROL – INDUSTRIAL USE ONLY

This product is a non-selective granular herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for non-cropland areas such as rights-of-way for pipeline, utility, highway, and railroad; and industrial areas such as airports, petroleum tank farms, lumberyards, and plant sites. 200 to 400 pound applications of this product per acre usually result in a non-productive condition of the soil for a period of a year or more. The duration of non-productivity is dependent upon rainfall, soil type, and other conditions.

This product should be applied as furnished with a seed spreader, a fertilizer spreader, a shaker-type applicator, or with any equipment which will distribute the chemical uniformly over the area to be treated. For control of annual and most perennial weeds and grasses, apply this product at the rate of 200 to 400 pounds per acre; on smaller plots, 0.5 to 1 pound per 100 square feet. Repeat spot treatment may be required when deep-rooted perennial weeds are present. To obtain best results, this product should be applied to the ground where it will be absorbed by the roots. Applications made early in the season have been found to give results superior to applications at a later seasonal date. This product may, however, be applied at any time of the year. Maximum effectiveness in arid regions is obtained when application is made just prior to the rainy season. Users should consult State Agricultural Experimental Stations or Extension Service weed specialists for recommendations as to use in their particular area.

**IMPORTANT (Not for Residential Use):** Do not apply on or near valuable woody or herbaceous plants or on areas where their roots may extend because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (or do not use same spreading equipment for other purposes). Do not use on croplands or any land to be used for subsequent cropping. Keep animals off treated areas.

Method

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

Rates

[field rates 0](#)

[field\\_rates 1](#)

- 

Timings

[early in the season](#)