

WEEDS CONTROL AT 0.125-0.25 OZ PER ACRE

General Information

This product is for use on Ornamental Turf, such as Lawns, Parks, Cemeteries, Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas. This product may also be used on Sod Farms.

USE PRECAUTIONS

Use lower rates for minimum chlorosis of the turf.

Do not apply this product to turf under stress from drought, insects, disease, cold temperatures, high temperatures of above 85°F on cool season grasses or poor fertility as injury may result.

Do not apply to turf less than 1 year old.

Do not use on Bahiagrass where it is the desired turf, as severe injury may result.

Do not plant ornamentals such as shrubs and trees in treated areas for at least 1 year after the last application or bedding plants for at least 2 years.

IMPORTANT

Addition of a nonionic surfactant of at least 80% active ingredient at 0.25 percent by volume (1 quart per 100 gallons) provides maximum performance, but may temporarily increase chlorosis of the turf.

Allow one week between the application of this product and other control (pesticide containing) products. (This guideline can be relaxed where severe insect or disease attack requires immediate treatment).

DO NOT USE ON FOOD OR FEED CROPS. Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply this product (except as instructed) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

When overseeding, wait 2 months (8 weeks) after application. Do not apply to any

body of water, including streams, irrigation water or wells. Do not apply where runoff water may flow onto agricultural land, as injury to crops may result.

Do not allow spray drift onto adjacent crops or other desirable plants or trees as injury may occur.

Follow these practices to minimize drift:

- Stop spraying if wind speed becomes excessive. Spray drift can occur at wind speeds less than 10 MPH. If sensitive plants are downwind, extreme caution must be used. Do not spray if winds are gusty.
- High temperatures, drought and low relative humidity increase the possibility of harmful spray drift. Caution must be used when these conditions are present and sensitive plants are nearby.
- Use large droplet size sprays to minimize drift.
- Use spray pressures of 35 psi or less when applying this product.

COMPATIBILITY

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent.

IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

RECOMMENDED SPRAYER CLEANUP

Immediately after spraying thoroughly remove all traces of this product from mixing and spray equipment as follows:

- 1) Drain tank, rinse interior surface of tank, then flush tank, boom and hoses with clean water for a minimum of 5 minutes.
- 2) Fill the tank with clean water, then add an ammonia cleaning solution. Use one gallon of ammonia (containing 3% active) per 100 gallons of water. Turn on sprayer long enough to flush through boom, hoses and nozzles. Stop spraying, but keep agitator working in the tank for 15 minutes, then drain.
- 3) Repeat Step 2.
- 4) Repeat Step 1.
- 5) Nozzles and screens should then be removed and cleaned separately. To remove traces of cleaning solution, rinse the tank thoroughly with clean water and flush through hoses and boom.
- 6) Flush boom and hoses with clean water for 5 minutes just prior to using the sprayer for the first time after the application of this product.

Limitations, Restrictions, and Exceptions

Use spray volumes of 20 to 90 gal/acre and pressures of 25 to 240 psi at use rate for weeds listed.

The required amount of this product should be added when the spray tank is half full of water and, with agitator running, add the proper amount of product. Finish adding the required amount of water. Continuous agitation is required to keep the product in suspension.

Spray preparations of this product may degrade in acid solutions if not used in 24 hours; it is stable in alkaline solutions. Thoroughly reagitrate before using.

Tank mixes with other registered herbicides should be tested for compatibility before full scale mixing. Use mechanical or bypass agitation to thoroughly mix the spray suspension. It is not necessary to premix this product with water in a separate container prior to adding it to the spray tank. This product should always be added to the tank first, before any other herbicides or adjuvants.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[N.A.](#)