

TURFGRASSES - AIDS IN SUPPRESSION (EXCEPT CALIFORNIA)

General Information

GENERAL PRODUCT INFORMATION

AFFIRM WDG is not for use on turf being grown for commercial seed production.

AFFIRM WDG is not for homeowner use to treat food crops.

AFFIRM WDG is for use in controlling or suppressing certain diseases in a Greenhouse, Lath or Shade house, Ornamentals (Field, Nursery, Container, and Residential and Commercial Landscapes), and Turf (including golf courses, sod, residential lawns, parks, athletic fields, and commercial and institutional grounds) composed of cool and warm season grasses such as Bluegrass, Bentgrass, Fescue, Ryegrass, Bermudagrass, St. Augustinegrass, Zoysiagrass, or their mixtures.

Apply AFFIRM WDG as a preventative or curative treatment in conjunction with good management practices.

RESISTANCE MANAGEMENT DIRECTIONS

AFFIRM WDG contains a Group 19 fungicide. Fungal isolates with acquired resistance to Group 19 may eventually dominate the fungal population if Group 19 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by AFFIRM WDG or other Group 19 Fungicides.

To delay fungicide resistance, consider:

- Avoiding the consecutive use of AFFIRM WDG or other target site of action Group 19 fungicides that have a similar target site of action, on the same pathogens.
- Using tank-mixtures or premixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Basing fungicide use on a comprehensive IPM program.

- Monitoring treated fungal populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

Limitations, Restrictions, and Exceptions

APPLICATION NOTES

- Repeat treatment on a 7-14 day schedule where environmental conditions are conducive to development of diseases. Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases.
- When symptoms are present, best control will be achieved by using a shorter interval.
- For best results apply after mowing.
- Apply to obtain thorough plant coverage in a minimum of 44 gallons water per acre.

Notes:

- With the exception for treatment of Fairy Ring, irrigation or rainfall soon after treatment with AFFIRM WDG will decrease disease control. For best control, do not irrigate for 12 hours after treatment.
- Use in alternation with fungicides that have different modes of action.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

4 hours

Timings

[After mowing.](#)