

PRETREATMENT OF NEW CONSTRUCTION

General Information

WITHIN FOOD AREAS AND NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

FOOD HANDLING ESTABLISHMENTS: Places other than private residences in which exposed food is held, processed, prepared or served.

Care should be taken to avoid depositing the product onto surfaces which are exposed or introducing the material into the air. Avoid contamination of food or food processing surfaces. APPLICATIONS OF THIS PRODUCT IN THE FOOD AREA OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

Apply in small amounts directly into cracks and crevices using a bellows duster or other suitable equipment capable of applying insecticide directly. Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas, when food is exposed and facility is in operation, also would be considered a food area.

Limitations, Restrictions, and Exceptions

PRETREATMENT OF NEW CONSTRUCTION (Preventive treatment for cockroaches and drywood termites): Apply powder to baseplates and plumbing voids at the rate of 4 pounds per 1,000 square feet. In attics, apply powder around perimeter of attics and around vent stacks at the rate of 4 pounds per 1,000 square feet.

Method

[N.A.](#)

Rates

[field rates 0](#)

-

Timings

[N.A.](#)