

POULTRY HOUSES-TO CONTROL DARKLING BEETLES

General Information

WITHIN FOOD AREAS AND NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

FOOD HANDLING ESTABLISHMENTS: Places other than private residences in which exposed food is held, processed, prepared or served.

Care should be taken to avoid depositing the product onto surfaces which are exposed or introducing the material into the air. Avoid contamination of food or food processing surfaces. APPLICATIONS OF THIS PRODUCT IN THE FOOD AREA OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

Apply in small amounts directly into cracks and crevices using a bellows duster or other suitable equipment capable of applying insecticide directly. Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas, when food is exposed and facility is in operation, also would be considered a food area.

Limitations, Restrictions, and Exceptions

POULTRY HOUSES-TO CONTROL DARKLING BEETLES: Remove birds before applying product. Apply at the rate of 1 to 2 pounds per 100 square feet over old litter or soil surface of the poultry house. Also apply to the posts, walls, framing and insulation of the poultry house. Do not apply directly to manure. Evenly spread fresh litter a minimum of four inches over all treated floor surfaces before introducing birds. Reapply after each grow-out period or after each sanitizing and cleaning of the poultry containment area in order to have good beetle control.

Method

[N.A.](#)

Rates

[field rates 0](#)

-

Timings

N.A.