

OUTDOOR USE - TELECOMMUNICATION TREATMENT

General Information

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

The product is a ready to use insecticide for use against the listed pests when used in accordance with the label directions.

For pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include (but are not limited to): industrial buildings and installations; houses; apartment buildings; laboratories; buses; greenhouses (non-commercial); mausoleums; sewers and underground tunnels; telecommunication conduits; utility vaults and conduits; and the nonfood/feed areas of stores, warehouses, vessels, railcars, trucks, trailers, aircraft (do not use in aircraft cabins), theaters, schools, nursing homes, hospitals, restaurants, hotels, and food manufacturing, processing, and servicing establishments.

The product is intended for applications with hand or power duster (Actisol DPA or similar equipment), with a paintbrush or by other suitable means, to hiding and runway areas and other places where insects are found.

To apply the product directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested area. Pay particular attention to cracks and crevices; service ducts; false floors and ceilings; wall voids; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets; refrigerators and sinks; around windows and doorframes; along baseboards; in attics and crawlspaces. The amount to be applied will vary with the site but should usually be in the range of 0.05 to 0.07 ounces per square yard (1.58 to 2.4 grams per square meter or 0.32 to 0.5 pounds per 1,000 square feet).

For indoor/outdoor uses: Repeat application only if there are signs of renewed insect

activity. Repeat application should be limited to no more than once per 7 days.

ATTENTION

Do not apply the product to edible crops.

Protect aquariums from dust.

Do not treat pets with the product.

During any application to overhead interior areas of structure, cover surface below with plastic shielding or similar material.

Do not apply in patient rooms while occupied or in any rooms while occupied by the elderly or infirm.

Do not apply to classrooms when in use.

Do not apply to institutions (including libraries, sports facilities, etc.) in the immediate area when occupants are present.

Do not use in warehouses where raw or cured tobacco is stored.

Do not use in warehouses while raw agricultural commodities for food or feed are being stored.

Do not use in greenhouses (non-commercial) where crops for food or feed are grown.

Limitations, Restrictions, and Exceptions

OUTDOOR USE

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Application to lawns, turf, and other vegetation;
- (3) Application to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to

impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crevice applications only.

Telecommunication Treatment

Apply the product into telecommunication conduits, cracks and crevices, junction boxes, etc. using suitable application equipment. The amount of the product to be applied will vary, but should usually be approximately 0.67 grams of dust per cubic foot of conduit.

The product can be used in areas showing signs of active infestations, such as underground telecommunication cable conduits, electrical cable conduits, junction boxes and manholes.

Applications should be made from both ends of a run of open conduit in cases where breakdown in the conduit is suspected. The product can also be applied directly to the interior of the conduits after cables have been installed. Apply at the rate of 0.67 grams of dust per cubic foot of conduit using equipment (i.e., Actisol DPA Duster) that is capable of delivering particles suspended in a low-pressure air system.

Method

[Spot treatment](#)

[Brush](#)

Rates

[field rates 0](#)

-

Timings

[Apply in areas showing signs of active infestation.](#)