

## **RAPESEED - GRASSHOPPERS**

Limitations, Restrictions, and Exceptions

**DIRECTIONS:** Use the designated amount of product in the following litres of water per hectare unless otherwise noted.

Ground Application: 200 to 1000 Litres of Water.

### **RECOMMENDED APPLICATIONS**

Use 575 mL - 1.25 L per hectare. Applications by air\* or ground equipment should begin when small larvae are present or when damage first appears. Two applications may be made. Do not apply later than 10 days before harvest.

\*FOR AIRCRAFT APPLICATION ON RAPESEED, USE NOT LESS THAN 10 LITRES OF WATER PER HECTARE. Refer to aerial application procedures.

**PRECAUTIONS:** Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

### **AERIAL APPLICATION PRECAUTIONS**

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of the label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on the label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

10 days

Rates

[field rates 0](#)

- 

Timings

[Begin when small larvae are present or when damage first appears.](#)