

CRANBERRY - PHYTOPHTHORA ROOT ROT

General Information

GENERAL INFORMATION

Ridomil Gold GR is a systemic fungicide which provides control of certain diseases caused by members of the Oomycete class of fungi on selected crops.

Note: Ridomil Gold GR is a systemic fungicide having a specific mode of action and could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive strains of fungi. If treatment is not effective following the use of Ridomil Gold GR as recommended, an insensitive resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a Ridomil Gold GR insensitive strain of fungi, neither Ridomil Gold GR nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance in your particular crop and disease control situation.

Do not use in nurseries, turf, or landscape plantings, unless specified in the label.

THE LABEL IS FOR FIELD USE ONLY AND IS NOT INTENDED FOR USE ON TRANSPLANT TRAYS, GREENHOUSES, LATH HOUSES, FLOAT HOUSES, HYDROPONIC PRODUCTION, OR IN BEDDING PLANT STRUCTURES.

Please refer to the label for the formula in calculating the amount of Ridomil Gold GR needed for band treatment.

Preharvest Intervals

Unless otherwise stated in the directions for use for a specific crop, there is no preharvest interval for Ridomil Gold GR.

Limitations, Restrictions, and Exceptions

CRANBERRY

For control of root rot of cranberries caused by Phytophthora spp.

Apply rate Ridomil Gold GR by ground or air. Make the first application in the fall after harvest. Make the second application in the spring, followed by a third application 45 days before harvest.

Notes: To avoid possible illegal residues, (1) Do not apply within 45 days before harvest, and (2) Do not apply more than 105 lbs./A per growing season.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

45 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[First application: In the fall after harvest.](#)

[Second Application: In the spring.](#)

[Third Application: 45 days before harvest.](#)