

TURFGRASS - CURATIVE TREATMENT

General Information

PRODUCT INFORMATION

Banol Fungicide is a completely water miscible fungicide concentrate. Banol Fungicide is intended for use on turfgrass or ornamental species which are subject to Pythium Blight (Cottony Blight, Grease Spot) or Phytophthora. It may be used after germination to control Pythium damping-off in turf, or on established turf to prevent or control Pythium Blight. Banol Fungicide does not afford broad spectrum disease control for Rhizoctonia sp. (brown patch), Sclerotinia sp. (dollarspot), or Helminthosporium sp. (leafspot). Use a broad spectrum fungicide if these diseases coexist with Pythium Blight.

Banol Fungicide is also labeled for prevention of root rot and damping-off caused by Pythium sp. and Phytophthora sp. at all stages of ornamental propagation.

COMPATIBILITY: Banol Fungicide is compatible with iprodione, ethazol, chloroneb, trichlorfon, triadimefon, chlorothalonil, diazinon, chlorpyrifos, and thiram. For control of other soil borne diseases such as Rhizoctonia sp. or Fusarium sp. Banol Fungicide can be tankmixed with thiophanate-methyl. Use the full rate of each fungicide and follow the most restrictive directions and restrictions appearing on the labels of these products.

USE RESTRICTIONS

Do not graze or feed clippings from treated areas to livestock.

Do not apply this product through any type of irrigation system.

Turfgrass and Ornamentals

Banol Fungicide is labeled for prevention of root rot and damping-off caused by Pythium sp. and Phytophthora sp. at all stages of turfgrass development and ornamental plant propagation, including seeding, transplanting, and potting.

Limitations, Restrictions, and Exceptions

Curative Treatment: 3-4 fl oz in 2-5 gallons of water

REMARKS

Do not apply more than a total of 12.25 fl oz (0.57 lb ai) of Banol Fungicide per 1,000 sq ft of turfgrass per year, equivalent to 25 lb ai/A/year.

ESTABLISHED TURF: Apply as a preventive treatment during periods of high temperature and humidity favorable to the development of Pythium Blight.

OVERSEEDDED AREAS: Apply after germination to control gallons of water. Pythium Damping-Off.

RETREATMENT: Retreat at 7-21 day intervals if conditions remain favorable for disease development. Use the lower rate for the shorter interval and the higher rate for the longer interval within the application rate range. Under severe disease pressure conditions, use the highest recommended rate and shortest interval.

RETREATMENT: Re-treat at 7-21 day intervals if conditions remain favorable for disease development. Use lower rate for the shorter interval and the higher rate for the longer interval within the application rate range. Under severe disease pressure conditions, use the highest labeled rate and shortest interval.

*NOT FOR USE ON SOD FARMS IN ARIZONA

Method

[Spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[Overseeded Areas: After germination.](#)