

ORNAMENTAL PLANTS - TRANSPLANTING (SEEDLINGS)

General Information

PRODUCT INFORMATION

Banol Fungicide is a completely water miscible fungicide concentrate. Banol Fungicide is intended for use on turfgrass or ornamental species which are subject to Pythium Blight (Cottony Blight, Grease Spot) or Phytophthora. It may be used after germination to control Pythium damping-off in turf, or on established turf to prevent or control Pythium Blight. Banol Fungicide does not afford broad spectrum disease control for Rhizoctonia sp. (brown patch), Sclerotinia sp. (dollarspot), or Helminthosporium sp. (leafspot). Use a broad spectrum fungicide if these diseases coexist with Pythium Blight.

Banol Fungicide is also labeled for prevention of root rot and damping-off caused by Pythium sp. and Phytophthora sp. at all stages of ornamental propagation.

COMPATIBILITY: Banol Fungicide is compatible with iprodione, ethazol, chloroneb, trichlorfon, triadimefon, chlorothalonil, diazinon, chlorpyrifos, and thiram. For control of other soil borne diseases such as Rhizoctonia sp. or Fusarium sp. Banol Fungicide can be tankmixed with thiophanate-methyl. Use the full rate of each fungicide and follow the most restrictive directions and restrictions appearing on the labels of these products.

USE RESTRICTIONS

Do not graze or feed clippings from treated areas to livestock.

Do not apply this product through any type of irrigation system.

Turfgrass and Ornamentals

Banol Fungicide is labeled for prevention of root rot and damping-off caused by Pythium sp. and Phytophthora sp. at all stages of turfgrass development and ornamental plant propagation, including seeding, transplanting, and potting.

Limitations, Restrictions, and Exceptions

Do not use for field grown ornamentals.

Banol Fungicide requires no agitation after initial mixing. Stock solutions of Banol Fungicide must be used within one week of mixing. Banol Fungicide does not cause

phytotoxic symptoms on the listed plants when used as directed. Trials on other ornamental species must be conducted before extensive use. Apply to commercially grown ornamentals in greenhouses and container grown ornamentals in nurseries by soil-directed drench only. Do not apply a soil drench to the foliage.

Use chemical resistant gloves when planting cuttings, seedlings, or seeds into the potting media that is still wet from a Banol Fungicide drench application. Do not apply more than 63 fl oz of Banol Fungicide (2.95 lb ai) per 1,000 sq ft per season.

When dipping plant roots, workers are required to use the stated personal protective equipment on this label and a chemical-resistant apron as a prudent measure.

REMARKS

Transplanting (Seedlings): Apply to the transplanting area using 3 quarts of solution/10 sq ft before transplanting. Do not make more than two applications per container per season. If a second application is needed, retreat at a minimum of 7-days to 6 months depending on the plant species and cultural practice.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[Pretransplant](#)