

## **COTTON - CHEMICAL HOE - 3-6 INCHES HEIGHT OF COTTON (AR, LA, MO, MS, TN, AND TX)**

### General Information

#### PRODUCT INFORMATION

Caparol 4L is a selective herbicide that may be applied either before or after weeds emerge for control of most annual broadleaf weeds and grasses, including groundcherry, lambsquarters, annual morningglory, malva, mustard, black nightshade, pigweed (carelessweed), purslane, Florida pusley, ragweed, smartweed, teaweed (prickly sida), barnyardgrass (watergrass), crabgrass, foxtail, goosegrass, junglerice, Panicum spp., signalgrass (and other Brachiaria spp.), and wild oats. Caparol 4L also controls shallow-germinating seedlings of cocklebur, coffeeweed, and sandbur. Caparol 4L will also provide partial control of spurred anoda (cottonweed), rough blackfoot (ironweed, cluster flaveria), and prairie sunflower in NM and western TX. Caparol 4L does not control johnsongrass, bermudagrass, other established perennials, or sprangletop at selective rates.

When applied before weeds emerge, Caparol 4L enters weeds through their roots. Thus, its effectiveness depends on moisture to move it into the soil. Under very dry soil conditions after application, a shallow cultivation or rotary hoeing will generally result in better weed control.

When applied to emerged weeds, Caparol 4L provides foliar knockdown and/or residual control of later germinating weeds, depending on the rate applied.

#### RESISTANCE MANAGEMENT

Caparol 4L Herbicide contains the active ingredient prometryn which inhibits photosynthesis at photosystem II (PSII, Site of Action Group 5). Some naturally occurring weed populations have been identified as resistant to Group 5 herbicides. Selection of resistant biotypes through repeated use of these herbicides or lower than labeled use rates in the same field, may result in weed control failures. A resistant biotype may be present where poor performance cannot be attributed to adverse environmental conditions or improper application methods. Contact your local Syngenta representative, retailer, crop advisor or extension agent to

determine if weeds resistant to this mode of action are present in your area. If resistant biotypes have been reported, use the full labeled rate of this product, apply at the labeled timing, and tank-mix with a different mode of action product so there are multiple effective modes of application for each suspected resistant weed.

## Principles of Herbicide Resistant Weed Management

### Scout and know your field

- Know weed species present in the field to be treated through scouting and field history. An understanding of weed biology is useful in designing a resistance management strategy. Ensure the weed management program will control all weeds present.
- Scout fields prior to application to determine species present and growth stage. Always apply this herbicide at the full labeled rate and correct timing for the weeds present in the field.

### Utilize non-herbicidal practices to add diversity

- Use diversified management tactics such as cover crops, mechanical weed control, harvest weed seed control, and crop rotation as appropriate.

### Use good agronomic practices, start clean and stay clean

- Use good agronomic practices that enhance crop competitiveness.
- Plant into weed-free fields utilizing tillage or an effective burndown herbicide for control of emerged weeds.
- Sanitize farm equipment to avoid spreading seed or vegetative propagules prior to leaving fields.

### Difficult to control weeds

- Fields with difficult to control weeds should be planted in rotation with crops that allow the use of herbicides with an alternative mode of action or different management practices.
- Difficult to control weeds may require sequential applications, such as a broad spectrum preemergence herbicide followed by one or more postemergence herbicide applications. Utilize herbicides containing different modes of action effective on the target weeds in sequential applications.

### Do not overuse the technology

- Do not use more than two applications of this or any other herbicide with the same mode of action in a single growing season unless mixed with an herbicide with a

different mode of action which provides overlapping spectrum for the difficult to control weeds.

Scout and inspect fields following application

- Prevent an influx of weeds into the field by controlling weeds in field borders.
- Scout fields after application to verify that the treatment was effective.
- Suspected- herbicide resistant weeds may be identified by these indicators
- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.
- Report non-performance of this product to your Syngenta retailer, Syngenta representative, or call 1-866-Syngent (866-796-4368). If resistance is suspected ensure weed escapes are controlled using an herbicide with an effective mode of action and/or use non-chemical means to prevent further seed production.

Prevent weed escapes before, during, and after harvest

Do not allow weed escapes to produce seed or vegetative structures such as tubers or stolons which contribute to spread and survival. Consider harvest weed seed management and control weeds post-harvest to prevent seed production.

## REPLANT AND ROTATIONAL CROPS

The following crops listed in the table in the label may be planted at the specified interval following application of Caparol 4L Herbicide to all crops on this label except cotton.

## ROTATIONAL USE RESTRICTIONS

- 1) DO NOT replant or rotate any crop in the table if more than 4 pt/A of Caparol 4L has been applied to the previous crop.
- 2) DO NOT use this table as guidance to replant or rotate any crop after treatment of cotton by Caparol 4L Herbicide.

Rotational Crop Restrictions Following Cotton

The following crops listed in the table in the label may be planted at the specified interval following application of Caparol 4L Herbicide to cotton.

## Precaution

- Cotton may be replanted in soil previously treated with Caparol 4L Herbicide. Application of a second preemergence treatment may result in crop injury.

## ROTATIONAL USE RESTRICTIONS

- 1) DO NOT replant or rotate any crop if more than one of the following applications of Caparol 4L Herbicide are used: pre-plant incorporated, pre-emergence or only one post-directed treatment.
- 2) DO NOT replant or rotate any crop until the following year where a lay-by or multiple applications of Caparol 4L Herbicide are made.

## APPLICATION PROCEDURES

Do not apply this product in a way that will make contact with workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Ground application (All uses): Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application.

Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 20 gal of spray mixture per acre for all preplant incorporated, preemergence, and postemergence applications (with or without surfactant) with ground equipment.

Use a pump with capacity to: (1) maintain nozzles at 35-40 psi, and (2) provide sufficient agitation in tank to keep mixture in suspension. A centrifugal pump which provides propeller shear action is recommended for dispersing and mixing this product. The pump should provide a minimum of 20 gal/minute/100 gal tank size circulated through a correctly positioned sparger tube or jets.

For preplant incorporated or preemergence application, use flat fan nozzle tips. For postemergence band application, use drop extraction tubes off-center nozzle tips. For postemergence broadcast application, use flat fan or off-center nozzle tips. Use flood nozzle tips only in AZ and CA for lay-by treatment in cotton at least 18 inches tall.

Use screens to protect the pump and to prevent nozzles from clogging. Screens

placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre using the formula given in the label.

Aerial application (Cotton and pigeon peas only): Use aerial application only where broadcast applications are specified. Use a minimum of 5 gal of spray mixture per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft above vegetation, using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Caparol 4L by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

## SPRAY EQUIPMENT

### Spray Drift Management

- Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.
- Apply only as a medium or coarser spray (ASABE standard 572.1) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.
- Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for ground boom application:

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Additional requirements for aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade

diameter.

- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.
- When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.
- Nozzles must always point backward parallel with air stream and never be pointed downward more than 45 degrees.

### Sensitive Areas

Apply the pesticide when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

### Seedbed Preparation

To ensure proper placement of Caparol 4L, seedbeds must be well prepared and as free as possible from trash and clods. A firm seedbed is best for obtaining effective weed control. Uniformity in height and width of seedbed is essential for proper postemergence applications of Caparol 4L. Beds should be low and flat. Take care to avoid planter marks. Wide planter packing wheels or rollers are recommended. Wheel furrows should be uniform in depth. Mount the sprayer so that it follows the same rows as the planter.

Refer to label for Tank Mix Information.

### Limitations, Restrictions, and Exceptions

#### COTTON

Restrictions: (1) Do not apply more than a total of 10.3 pt of Caparol 4L on sandy loam soil or 11.9 pt of Caparol 4L on medium or fine soil per acre per year. (2) Do not feed treated forage to livestock or graze treated areas. (3) Do not use on glandless cotton varieties.

#### Caparol 4L Alone

Caparol 4L may be applied preplant incorporated or preemergence and/or

postemergence as specified in the following tables. The postemergence applications may follow preplant incorporated or preemergence treatments of Caparol 4L.

### Postemergence-Directed

Be especially careful when applying Caparol 4L postemergence to prevent contact of the spray with cotton leaves, or injury may occur. Use precision application equipment so the spray is accurately directed to the base of the cotton plants and still thoroughly covers the soil and weeds beneath the cotton plants. Apply during calm periods to prevent drift. Use leaf lifters or shields if leaf contact cannot be avoided merely by directing the spray. Apply only when all cotton plants have exceeded the minimum recommended height shown in Tables 3 and 4. Apply to level, well prepared surfaces such as relatively clod-free beds made with bed-shapers.

When applying to emerged weeds, add 2 qt of surfactant per 100 gal of spray mixture. Use a surfactant that is compatible with Caparol 4L when applied in cotton and is approved by EPA for use on food and feed crops. Examples include X-77, Tronic, and Triton.

Precautions: (1) Application to furrow-planted cotton before furrows are leveled (plowed in) may cause crop injury. (2) Application to cotton under stress from drought, cultivator damage, or fertilizer application may cause crop injury.

Chemical Hoe (Emerged Weeds only): Apply Caparol 4L at the appropriate rate in Table 3, two or three times if necessary. In cotton 3-6 inches tall, be extremely careful to avoid spray contact with cotton leaves by applying Caparol 4L with a precision applicator equipped with fenders or shields, such as Bell Row Shield, Dickey Fenders, or W&A Fenders. In cotton less than 10 inches tall, apply only if cotton is bed or flat-planted.

Height of Weeds: less than 1 inch

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Directed](#)

Rates

## [field\\_rates 0](#)

- 

Restricted Entry Interval

12 hours

Exception: If the product is soilinjected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Postemergence \(Weed\)](#)