

## **SURFACE FEEDING INSECTS**

### General Information

#### RESTRICTIONS

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Aerial application is prohibited.
- Use this product only on the turf sites permitted by this label.
- Do not apply this product using hand-held or chest-mounted spreaders.
- The applicator must water-in the product after application, or notify the property owner that he or she must water-in after application on the same day.
- Limit the number of applications to turf to 3 per calendar year with minimum retreatment intervals of 7 days.
- Limit the total of all applications to 24.5 lbs of active ingredient (395 lbs of product) per acre per year for controlling mole crickets and grubs and to 16.2 lbs of active ingredient (261 lbs of product) per acre per year for controlling surface feeding insects.
- Apply this product directly to the lawn or garden area. Water treated area as directed on this label.
- Do not water to the point of run-off. Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

#### Additional Restrictions For Turf on Golf Courses

- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams,

marshes, natural ponds, or estuaries.

- Broadcast use is limited to tees and greens. Use on fairways is limited to spot treatments.
- Post-application watering-in is mandatory.

#### PRODUCT SPECIFICATIONS

DYLOX 6.2 - Granular Insecticide containing 6.2% trichlorfon

GRANULAR INSECTICIDE CONTROLS - White Grub Larvae (such as larvae of Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, Aphodius spp., May or June beetles (Phyllophaga spp.), northern and southern masked chafers (Cyclocephala spp.), oriental beetle and sugarcane grubs (Tomarus spp.))

- Mole crickets
- Annual bluegrass weevil
- Billbug larvae
- Sod webworms and cranberry girdler
- Cutworms and armyworms
- Chinch bugs

WHERE TO APPLY - All types of turf where pests have been seen or found excluding turf where application is not allowed. See IMPORTANT: above.

COVERAGE - One bag (30 lbs) will treat 10,000 sq ft for white grubs, annual bluegrass weevil, chinch bugs, cranberry girdler, and mole crickets.

- One bag (30 lbs) will treat 15,000 sq ft for sod webworms, armyworms, and cutworms.

HOW TO APPLY - Do not apply with hand-held or chest-mounted spreader.

- Apply granules using a broadcast or drop type spreader.

QUESTIONS For questions or comments, call toll free 1 (800) 331-2867.

Limitations, Restrictions, and Exceptions

#### SURFACE FEEDING INSECTS

Refer to chart for spreader settings (Make sure your spreader is calibrated for this

product before application).

### How to Apply

- For properly managed turf remove excessive thatch, which encourages insects. Rake or de-thatch the area and remove the debris to expose insect hiding places.
- Water the turf and cut to proper height.
- Apply granules with a broadcast or drop type spreader.
- To release the active ingredient from the granule, the applicator must water-in after application or notify the property owner that he or she must water-in, on the same day as application, with sufficient water to thoroughly wet the remaining thatch. If sufficient rainfall to thoroughly wet the thatch occurs after application but before watering-in, additional watering-in is not required.

When to Apply - Treat when the insects are present.

- To control multiple generations of sod webworm larvae, additional treatments at 30- to 40-day intervals during the summer months may be required. Do not make more than 3 applications per calendar year.

### Additional Restrictions For Turf on Golf Courses

- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, or estuaries.
- Broadcast use is limited to tees and greens. Use on fairways is limited to spot treatments.
- Post-application watering-in is mandatory.

Apply Dylox 6.2 Granular Insecticide uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply Dylox 6.2 Granular Insecticide. Do not use spreaders that would apply the material in concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

Method

[Spot treatment](#)

[Broadcast](#)

Rates

[field rates 0](#)

- 

Timings

[Treat when the insects are present. Repeat as needed.](#)