

LOW VOLUME BASAL BARK TREATMENTS

General Information

GENERAL INFORMATION

Stalker is an aqueous formulation that is readily mixable with water, diesel oil, or recommended seed oils and penetrating oils. Stalker is to be mixed with water or a penetrating oil and applied as a spray to cut stumps and frilling cuts for the control of brush. Stalker should be mixed with a penetrating oil for application to the basal area of brush and trees. Adequate agitation should be maintained with all Stalker emulsion mixtures to prevent phase separation. Prior to actual tank mixing with other products, herbicides and carrier oils, compatibility testing in small containers is recommended!

An application of Stalker is recommended for control of brush in grass pasture and rangeland and noncropland areas such as railroad, utility, highway, and pipeline rights-of-way, utility plant sites, petroleum tank farms, pumping installations, fence rows, storage areas, non-irrigation ditchbanks including grazed or hayed areas within these sites. Stalker is also recommended for use with asphalt and asphalt slurries to control weeds on road shoulders, under pavement, in roadside cracks and crevices, and to prevent weed encroachment on highways and paved surfaces.

Stalker is also recommended for control of undesirable vegetation along forest roads, non-irrigation ditchbanks, and the establishment and maintenance of wildlife openings.

Limitations, Restrictions, and Exceptions

LOW VOLUME BASAL BARK TREATMENTS

Mix 8.0-12.0 fluid ounces of Stalker in one gallon of diesel oil or a penetrating oil. To control mixed brush species with up to 4 inch stem diameter at breast height, spray to wet the lower 12-18 inches of the stem with the Stalker oil mixture (include the root collar area). DO NOT over apply causing dripping or puddling. Maintain uniform mixtures with frequent agitation. Avoid application on sites that have been mowed prior to application resulting in a high density of stump resprouts containing multiple, small (1/2 inch diameter or less) stems. Application sites containing high

stem densities and multiple, small (1/2 inch diameter or less) stems should be foliar treated with low volume backpack or fixed boom applications. Stalker may be tank-mixed with Garlon 4, Brush Killer 800 and other basal products to broaden the spectrum of control. Consult the herbicide labels for rates and susceptible brush species. When tank-mixing, follow all precautions on the tank-mix product label and always follow the most restrictive label. Use a tank mix of 3 to 5% Stalker plus 15 to 20% Garlon 4 in basal oil to control black locust, honey locust, hackberry, elms and other species listed on manufacturer's labels. Use the higher rate of Stalker (5%) in areas containing sassafras, oak, hickory, cherry, and maples or in the southern 2/3's of the U.S. A tank-mix of 3% Stalker herbicide+ Garlon 4 herbicide is effective in the Northeastern U.S.

Method

[Broadcast/Foliar Ground](#)

[Foliar spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)