

LOW VOLUME FOLIAR APPLICATIONS

General Information

GENERAL INFORMATION

Stalker is an aqueous formulation that is readily mixable with water, diesel oil, or recommended seed oils and penetrating oils. Stalker is to be mixed with water or a penetrating oil and applied as a spray to cut stumps and frilling cuts for the control of brush. Stalker should be mixed with a penetrating oil for application to the basal area of brush and trees. Adequate agitation should be maintained with all Stalker emulsion mixtures to prevent phase separation. Prior to actual tank mixing with other products, herbicides and carrier oils, compatibility testing in small containers is recommended!

An application of Stalker is recommended for control of brush in grass pasture and rangeland and noncropland areas such as railroad, utility, highway, and pipeline rights-of-way, utility plant sites, petroleum tank farms, pumping installations, fence rows, storage areas, non-irrigation ditchbanks including grazed or hayed areas within these sites. Stalker is also recommended for use with asphalt and asphalt slurries to control weeds on road shoulders, under pavement, in roadside cracks and crevices, and to prevent weed encroachment on highways and paved surfaces.

Stalker is also recommended for control of undesirable vegetation along forest roads, non-irrigation ditchbanks, and the establishment and maintenance of wildlife openings.

Limitations, Restrictions, and Exceptions

LOW VOLUME FOLIAR APPLICATIONS

Stalker may be applied as a low volume foliar application. Mix 3-5% Stalker in water and adjuvant or in a penetrating oil. For small brush spray down on the crown to cover approximately 70% of the plant foliage. For larger brush insure coverage on as much of the crown as possible and spray at least two sides of the plant. Stalker may be tank-mixed with other labeled herbicides. Use a tank mix of 3 to 5% Stalker plus 15 to 20% Garlon 4 in basal oil to control black locust, honey locust, hackberry, elms and other species listed on manufacturer's labels. Use the higher rate of

Stalker (5%) in areas containing sassafras, oak, hickory, cherry, and maples or in the southern 2/3's of the U.S. A tank-mix of 3% Stalker + Garlon 4 is effective in the Northeastern U.S.

Refer in the label for the spray solution mixing guide for low volume foliar applications.

INVERT EMULSIONS

Stalker can be applied as an invert emulsion carrier. The carrier is a thick invert water-in-oil spray emulsion designed to minimize spray drift and spray run-off, resulting in more herbicide on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions. DO NOT exceed 3 quarts/Acre of Stalker herbicide.

Method

[Broadcast/Foliar Ground](#)

[Foliar spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[N.A.](#)