

# **CONIFERS AND TREES IN COMMERCIAL NURSERIES, ETC. - TRUNK INJECTION**

## General Information

### GENERAL APPLICATION INSTRUCTIONS

Apply FOSPHITE by various application methods, including foliar sprays, trunk injection, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply FOSPHITE with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

### LOW VOLUME AND ULTRA-LOW VOLUME APPLICATIONS

Label rates are written for conventional application equipment. For LV/ULV equipment, use per acre rate of FOSPHITE and adjust amount of water to equipment needs. Always conduct a phytotoxicity test on a small area before applying to a large area in order to assess any potential risk to plants.

### GENERAL INSTRUCTIONS FOR APPLICATIONS FOOD CROPS

FOSPHITE fungicide is used for control of Downy Mildew, Powdery Mildew, *Erwinia amylovora*, *Pythium*, *Fusarium*, *Rhizoctonia* and *Phytophthora* disease (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/foliar rot, fire blight, heart rot, pink rot, *pythium* leak, red stele, leather rot, etc.) of agronomic crops. (See specific crop groupings for additional diseases.) This product may be applied up to the day of harvest.

### Limitations, Restrictions, and Exceptions

FOSPHITE applied in conjunction with good cultural management practices will control *Phytophthora* root rot of conifers including (but not limited to): Douglas Fir, spruce and pines also other *Phytophthora* disease of Oaks (Sudden Oak Death) and forest trees. Use the higher rates when disease pressure is severe. Do not apply to trees that are under heat or moisture stress.

### Application Program

Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.

Rate: 3 teaspoons / linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliters/linear meter of canopy width or 5 centimeters DBH.)

Inject directly in to tree.

Note: Do not graze livestock in treated conifer nurseries and/or plantations. Do not feed forage to livestock from treated nurseries and/or plantations.

Method

[Injection](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

4 hours

Timings

[N. A.](#)