

# **FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS**

## General Information

### GENERAL INFORMATION ON THE USE OF THIS PRODUCT

For use on plants intended only for aesthetic purposes or climatic modifications and being grown in interior plantscapes, ornamental gardens or parks, or lawns and grounds. Not for use on plants grown for sale or other commercial use, or for commercial seed production or research purposes.

### GENERAL APPLICATION INSTRUCTIONS

This product formulation mixes readily with water and other aqueous carriers, and controls a wide spectrum of insects and mites on trees, shrubs, foliage plants, non-bearing fruit and nut trees, and flowers in interiorscapes including hotels, shopping malls, office buildings, etc., and outdoor plantscapes, such as around residential dwellings, parks, institutional buildings, recreational areas, athletic fields and home lawns. Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application.

This product may be tank-mixed with other pesticides, including insect growth regulators. When tank mixing this product with other pesticides, observe all precautions and limitations on each separate product label. The physical compatibility of this product may vary with different sources of pesticide products, and local cultural practices. Any tank mixture which has not been previously tested should be prepared on a small scale (pint or quart jar), using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.

### RESISTANCE MANAGEMENT

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area.

Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products

with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

#### FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS

Applications of this product are permitted in both food/feed and nonfood areas of food/feed handling establishments as a general surface, spot, or crack and crevice treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

Permitted use sites also include, but are not limited to: Aircraft (Do not use in aircraft cabins), food manufacturing plants, grain bins.

**General Surface Application:** Do not use this application method in food/feed handling establishments when the facility is in operation or foods/feeds are exposed. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias, and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean fresh water.

**Spot, Crack and Crevice Application:** Spot or crack and crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food.

#### Foam Applications

This product may be converted to foam and used to treat structural voids. Dilute 0.33 to 1.0 fluid oz. of this product per gallon of water and add the manufacturer's recommended volume of foaming agent to produce a 0.02% to 0.06% foam

concentration. Verify before treatment that the foaming agent is compatible with this product.

Method

[Spot treatment](#)

[Surface](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

Timings

[N.A.](#)