

# **ORNAMENTAL PLANTS AND NON-BEARING FRUIT AND NUT TREES AND VINES - CONTROL OF SNAIL AND SLUG**

## General Information

### GENERAL PRECAUTIONS AND RESTRICTIONS

Do not use on crops for food or forage. Use only according to label directions. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

FOR PEST CONTROL: Mesurol 75-W effectively controls certain specified destructive pests on ornamentals. Use specified quantities of Mesurol 75-W in sufficient water to give complete coverage of foliage. The type of equipment used will determine the concentration required. Thorough coverage is necessary for maximum control.

The product mixes readily with water to form a suspension usable in all conventional spray equipment. Maintain adequate agitation in the spray tank to keep the wettable powder in suspension. Do not apply with oil.

Schedule applications in accordance with local conditions.

Performance may be reduced when the spray solution has a pH greater than 7. The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum activity.

### RESISTANCE MANAGEMENT STATEMENT

Repeated use of pesticides with similar modes of action may lead to a buildup of resistant pest strains. For optimum resistance management, Gowan Company recommends rotating or mixing products from different classes based on mode of action.

### Limitations, Restrictions, and Exceptions

### ORNAMENTAL PLANTS AND NON-BEARING FRUIT AND NUT TREES AND VINES

- Grows in Nurseries and Greenhouses and Mature Ornamental Plantings

Foliar Application: To control pests on foliage, flowers, and other ground plant parts, spray plants and foliage until thoroughly wet. Do not make more than 2 applications per year per crop. Applications must be at least 10 days apart.

- Use extreme caution when applying in conjunction with foliar fertilizer applications.
- Before treating a large number of ornamental plants with Mesurol alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7-10 days prior to treating large areas to reduce the possibility of plant injury.
- Do not make more than 2 applications per year per crop. Applications must be at least 10 days apart.

Method

[Foliar spray](#)

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

24hours

Timings

[N.A.](#)