

# **ORNAMENTAL LAWNS AND TURF - BENTGRASS TURF**

## General Information

### GENERAL INFORMATION

TRIAMINE II is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas, woody plants, along Highways, Rights-of-Way, and other similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### USE PRECAUTIONS

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not spray on carpetgrass, dichondra, nor on grasses where desirable clovers are present. Except as noted, use a lawn type sprayer with coarse spray as wind drift is less likely. Avoid fine mists. Do not spray exposed roots of desirable plants. Maximum control of weeds will be obtained by applying in spring and early fall when weeds are actively growing. Use the higher rate for hard-to-kill weeds. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) Turf should not be mowed 1 to 2 days before and following application; (2) Reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in injury.

Triamine II can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior

to mixing in the application equipment. All regulations, either state or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

## Limitations, Restrictions, and Exceptions

### ORNAMENTAL LAWNS AND TURF - BENTGRASS TURF

- On closely mowed Bentgrass, other than golf course greens and tees, apply Triamine II at a maximum rate of 0.67 ounce in 3 gallons water per 1,000 square feet (1.75 pints per acre) when weeds are actively growing. Exercise care to avoid overdosing Bentgrass or injury may result. Slight turf yellowing should disappear after about 1 week.

Low Volume Applicator: When using a Low Volume Applicator (such as Lawn Doctor's "Turf Tamer") use Triamine II at a rate of 3 to 4 pints per acre (1.1 oz. to 1.5 oz. in 1/2 gallon of water per 1,000 sq. ft.).

#### Method

[Spray](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

•

#### Timings

[May or mid-August through September.](#)