

## **VEGETABLE CROPS - CABBAGE - APHIDS, ETC.**

### General Information

#### GENERAL INFORMATION

Diazinon AG500 is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use in greenhouses. Do not use in barns or dwellings. Do not use on livestock. Aerial application is prohibited, except as allowed on the label for use on lettuce.

In the directions for use which follow, best control is obtained when application is made when infestation is noticed.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Do not use water from treated cranberry/watercress beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.

Note: The California Department of Pesticide Regulation has set a 5-day reentry period for diazinon on peaches and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

Failure to follow the directions for use and precautions on the label may result in poor insect control, crop injury, or illegal residues.

Refer to the label for tank mix information.

#### Limitations, Restrictions, and Exceptions

#### VEGETABLE CROPS

For foliar or soil applications, DIAZINON AG500 should always be applied as a spray

in sufficient water to assure thorough coverage of the foliage or soil. DIAZINON AG500 may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Notes: 1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see TIMING/SPECIAL DIRECTIONS) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches; for subterranean cutworms, incorporate 3-6 inches. 2) To Protect Bees: Foliar applications of this product should not be made to corn during the pollen shed period.

CABBAGE:Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers - Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. NOTE: Diazinon will not control organophosphate-resistant leafminers.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Soil incorporation](#)

Pre-Harvest Interval

21 days

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[Apply as insects occur.](#)