

## **SUGARCANE - LOUISIANA**

### General Information

#### GENERAL INFORMATION

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane, and certain other crops specified on the label. It is also effective in industrial sites for control of most annual and many perennial broadleaf and grass weeds. This product may be applied before or after weeds emerge.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on the label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your state Agricultural Extension Service for specific recommendations.

Since this product acts mainly through root absorption, its effectiveness depends on moisture to move it into the root zone. If weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is nonflammable.

Avoid using near adjacent desirable plants or in greenhouses, or injury may occur.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap as crop injury may result.

Where the use directions give a range of rates, use the lower rate on coarse-textured soil and soil low in organic matter; use the higher rate on fine-textured soil and soil high in organic matter.

NOTE: SIPCAM AGRO USA, INC. does not recommend applications in combination with other herbicides or oils, except as specifically described on the label.

## APPLICATION PROCEDURES

**GROUND APPLICATION:** Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gals. of spray mixture/A for all preplant incorporated, preplant surface, preemergence, and postemergence applications (with or without oil or surfactant) with ground equipment.

**AERIAL APPLICATION:** Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 qt. of water for each quart of this product applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gals. of water/A. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

**APPLICATION IN WATER OR LIQUID FERTILIZER:** Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence, preplant incorporated, or preplant surface ground application on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, or crop injury may occur.

Refer to the label for directions on band applications.

## Limitations, Restrictions, and Exceptions

### SUGARCANE

For control of annual weeds during the summer fallow period, apply rate to weed-free beds. Follow normal weed control after planting.

**USE PRECAUTIONS AND RESTRICTIONS:** To avoid crop injury if making a 4 pts./A application during the summer fallow period, do not exceed 8 qts./A during the remainder of the growing season, or illegal residues may result.

### Method

[Broadcast/Foliar Air](#)

## Broadcast/Foliar Ground

Rates

[field rates 0](#)

[field rates 1](#)

- 

Restricted Entry Interval

12 hours

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Apply immediately after bed formation.](#)