

PECAN - FALL WEBWORM, PECAN NUT CASEBEARER, PHYLLOXERA, SPITTLEBUG

General Information

GENERAL INFORMATION

Lorsban 50W insecticide in water soluble packets is designed for use as a cover or foliar spray to control various pests. The pests controlled are listed in the accompanying table. Generally, Lorsban 50W is compatible with insecticides, miticides, and fungicides commonly recommended for agricultural crops except for alkaline materials such as Bordeaux mixture and lime. Prepare a small amount of the spray mixture and check for compatibility before mixing a large volume of spray.

General Use Precautions and Restrictions

- Insect control may be reduced at low spray volumes under high temperature and wind conditions.
- Some reduction in insect control may occur under unusually cool conditions.
- Flood Irrigation: To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours following application of Lorsban 50W.
- Chemigation: Do not apply this product through any type of irrigation system.

Handling Precautions for Water Soluble Packets: Do not remove water soluble packet from the overpack except for immediate use. Do not allow packets to come into contact with water prior to use. Do not handle water soluble packets with wet hands or wet gloves. Do not open water soluble packets. Partial use of packets is not allowed. Carefully reseal package containing unopened water soluble packets and protect package from moisture.

Application Guidelines

Broadcast Foliar Application

Apply with conventional power-operated spray equipment using nozzles and spray pressures recommended for insecticides. Apply Lorsban 50W in a spray volume of

not less than 2 gallons per acre (gpa) for aerial application equipment (fixed wing or helicopter) or not less than 10 gpa for ground equipment, unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy.

Ground Application: Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's recommendations for insecticide nozzles with respect to nozzle type, pressure, and spacing.

Broadcast Soil Application

Apply with conventional power-operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets (235-400 microns). Unless otherwise indicated, a spray volume of 10 gpa or more is recommended. For band application, use proportionally less spray volume.

Aerial Application

Use a minimum spray volume of 2 gpa and follow recommendations for best management practices for aerial application, above.

Marking of swaths by flagging, permanent markers or use of GPS equipment is recommended.

Limitations, Restrictions, and Exceptions

PECAN

- Apply 1 lbs/100 gallons (1 1-lb packet).
- For control of Fall webworm, Pecan nut casebearer, spittle bug and Phylloxera.
- For best Phylloxera control, make 2 applications at 7 to 10 day interval starting at bud swell and ending by early leaf expansion.
- Apply Lorsban 50W as a foliar spray at the dosages indicated to control pests. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop and apply as a concentrate or dilute spray using conventional, power operated spray equipment. For dilute sprays applied to tree nut

crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of Lorsban 50W per acre. Treat when pests appear or in accordance with local conditions. Aerial application may result in less effective insect control because of reduced coverage. Consult your State agricultural experiment station, certified Pest control advisor, or Extension service specialist for specific use information in your area.

- REI is 4 hours to fruit trees and 24 hours to nut trees.

Specific Use Precautions:

- Lorsban 50W is highly toxic to bees exposed to direct treatment and should not be applied when bees are actively foraging in the treated area.
- To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours of application of Lorsban 75WG.

Specific Use Restrictions:

- Do not make a second application of Lorsban 75WG or other product containing chlorpyrifos within 10 days of the first application.
- Do not allow meat or dairy animals to graze in treated orchards.

Method

[Foliar spray](#)

Pre-Harvest Interval

28 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

24

Timings

[When pests appear.](#)

[Phylloxera : At bud swell.](#)