

WHEAT, BARLEY AND RYE (NOT UNDERSEEDED WITH LEGUMES) - POSTEMERGENCE - ANNUAL AND BIENNIAL WEEDS CONTROL

General Information

GENERAL INFORMATION

Nufarm WEEDONE LV4 Solventless Broadleaf Herbicide contains the isooctyl (2-ethylhexyl) ester of 2,4-D, the original and one of the best low volatile esters. In cropland, this herbicide is more effective than amines for controlling hard-to-kill weeds such as bindweed, thistles, smartweeds, wild garlic, curled dock, tansy ragwort and wild onions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

Limitations, Restrictions, and Exceptions

WHEAT, BARLEY and RYE (not underseeded with legumes)

DIRECTION

Do not spray grain in boot to dough stage.

0.5 to 2 pints / acre: Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds

are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage. Do not apply to grain in the seedling stage.

NOTE

Control Mustards (except blue mustard) and Pepperweeds (except perennial).

These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for application: beggarticks; common broomweed; knotweed; mallow (venice, dwarf, little); prickly lettuce; Russian thistle; smartweeds (annual species); vervains; wild carrot.

Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

RESTRICTIONS AND LIMITATIONS FOR USE ON CEREAL GRAINS

Do not mix with oil for crop uses.

Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.

Do not feed treated straw to livestock if a preharvest treatment as described above is applied.

- For the tank mixtures for cereal grains refer in the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[After grain is fully tillered \(usually about 4 to 8 inches high\) but not forming joints in](#)

the stem.