

FIELD CORN - PREEMERGENCE - BROADLEAF WEEDS CONTROL OR SUPPRESSION

General Information

GENERAL INFORMATION

Nufarm WEEDONE LV4 Solventless Broadleaf Herbicide contains the isooctyl (2-ethylhexyl) ester of 2,4-D, the original and one of the best low volatile esters. In cropland, this herbicide is more effective than amines for controlling hard-to-kill weeds such as bindweed, thistles, smartweeds, wild garlic, curled dock, tansy ragwort and wild onions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

Limitations, Restrictions, and Exceptions

FIELD CORN - Preemergence

DIRECTIONS

To control broadleaf weeds and suppress annual grasses, apply 3 to 5 days after planting, but before corn emerges. Use high rate on soil high in organic matter. Do not use on light, sandy soil or where soil moisture is inadequate for normal weed growth.

NOTE

Control Mustards (except blue mustard) and Pepperweeds (except perennial).

These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for application: beggarticks; common broomweed; knotweed; mallow (venice, dwarf, little); prickly lettuce; Russian thistle; smartweeds (annual species); vervains; wild carrot; alfalfa; bindweed (hedge, field & European); Canada thistle; docks; dogbanes; goldenrod; ground ivy; hawkweed (orange); hoary cress; many-flowered aster; nettles (including stinging); tansy ragwort; vervains; western ironweed; wild garlic; wild onion.

Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

RESTRICTIONS AND LIMITATIONS FOR USE ON FIELD CORN

Do not forage or feed corn fodder for 7 days following application.

Note: Hybrids vary in response to 2,4-D and some are easily injured. Contact seed company or your Agricultural Experiment Station or Extensional weed specialists for this information.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Preemergence \(Crop\)](#)