

# **LANDSCAPE ORNAMENTALS, ETC. - ANNUAL BLUEGRASS WEEVIL LARVAE**

## General Information

### DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### RESTRICTIONS

- Use this product only as directed by its label. Do not apply this product to sites not listed in the APPLICATIONS table.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Aerial application is prohibited.
- Do not apply this product through any type of chemigation system.
- Do not use this product on turf being grown for sale or other commercial use as sod. Do not use this product on turf being grown for commercial seed production or for research purposes except for research involving use of this product that is consistent with this label.
- Do not apply this product to crops grown for food or forage.
- Do not make foliar applications to trees, shrubs, bushes, or flowers. Ornamental plants may be treated only by applying spray to the soil at the base of the plant.
- For applications to turf, limit the number of applications to 3 per calendar year with minimum re-treatment intervals of 7 days.
- Do not apply more than 24.5 lbs of active ingredient per acre per year for controlling grubs and mole crickets. Do not apply more than 16.2 lbs of active ingredient per acre per year for controlling surface feeding insects.
- Post-application irrigation is required.
- Do not apply when turf grass areas are water-logged or the soil is saturated with water (i.e. will not allow penetration into the root zone of the plant). Do not apply to the point of run-off or puddling of irrigation water following application.
- Do not use this product on plants being grown for sale or other commercial use.

### Limitations, Restrictions, and Exceptions

## APPLICATIONS

- Mix specified dosage of DYLOX 420 SL Turf and Ornamental Insecticide in sufficient water to provide adequate distribution during application to lawn, turf, or groundcover area. To ensure that the active ingredient reaches the soil surface, apply with a minimum of 2 gal of water per 1,000 sq ft.
- Minimize the amount of thatch present prior to application. After application (on the same day), irrigate with sufficient water to thoroughly wet the remaining thatch and top layers of the soil. If sufficient rainfall to thoroughly wet the thatch and top layers of soil occurs after application and prior to irrigation, further irrigation is not required. Under dry conditions, pre-water prior to application for optimum control of white grubs and weevil larvae.
- If applied to newly mowed *Poa annua* irrigate as soon as possible after application, particularly under conditions of high temperature and humidity. Potential injury may be exacerbated by tank mixing with other products, or by the addition of surfactants or other additives.
- If the turf is being maintained at a mowing height of greater than 1 inch, then the maximum specified application rate (6.9 fl oz per 1,000 sq ft) may be required during periods of high pest pressure.
- Do not use DYLOX 420 SL Turf and Ornamental Insecticide in hose-proportioned sprayers.
- Note: Do not use treated area or clippings from treated areas for feed or forage.
- For ornamentals, apply to soil around the plant.

### Method

[Broadcast/Foliar Ground](#)

[Surface](#)

### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

### Timings

[N.A.](#)