

# **NARCISSUS - BULB FLY**

## General Information

### DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### RESTRICTIONS

- Use this product only as directed by its label. Do not apply this product to sites not listed in the APPLICATIONS table.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Aerial application is prohibited.
- Do not apply this product through any type of chemigation system.
- Do not use this product on turf being grown for sale or other commercial use as sod. Do not use this product on turf being grown for commercial seed production or for research purposes except for research involving use of this product that is consistent with this label.
- Do not apply this product to crops grown for food or forage.
- Do not make foliar applications to trees, shrubs, bushes, or flowers. Ornamental plants may be treated only by applying spray to the soil at the base of the plant.
- For applications to turf, limit the number of applications to 3 per calendar year with minimum re-treatment intervals of 7 days.
- Do not apply more than 24.5 lbs of active ingredient per acre per year for controlling grubs and mole crickets. Do not apply more than 16.2 lbs of active ingredient per acre per year for controlling surface feeding insects.
- Post-application irrigation is required.
- Do not apply when turf grass areas are water-logged or the soil is saturated with water (i.e. will not allow penetration into the root zone of the plant). Do not apply to the point of run-off or puddling of irrigation water following application.
- Do not use this product on plants being grown for sale or other commercial use.

### Limitations, Restrictions, and Exceptions

## NARCISSUS - BULB FLY

- Apply specified dosage in 100 gallons of water as a drench per 1,000 feet of row. Direct stream to base of plants at beginning of adult fly activity (early May to June). Repeat treatments annually.

Method

[Drench](#)

Rates

[field\\_rates 0](#)

- 

Timings

[At beginning of adult fly activity \(early May to June\).](#)