

CANOLA, CRAMBE AND RAPESEED - APHID, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zetacypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zetacypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

Limitations, Restrictions, and Exceptions

CANOLA, CRAMBE AND RAPESEED

RESTRICTIONS

Application Interval: Do not make applications less than 14 days apart.

Maximum Amount per Application: Do not apply more than 5.95 ounces of product or 0.053 pound active ingredient per application.

Maximum Amount of Hero EW Insecticide allowed per Season: Do not apply more than 11.9 ounces of product or 0.106 pound active ingredient per acre per season.

Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

Remarks

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).

Armyworm Fall: Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.

Diamondback moth: Pyrethroid resistance is common for these pests. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer to the resistance management statement in the DIRECTION FOR USE section of the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

35 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)