

BUSHES, VINES AND TREES - CANEBERRIES - TWOSPOTTED MITE, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zeta-cypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zeta-cypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

Limitations, Restrictions, and Exceptions

CANEBERRIES

RESTRICTIONS

Application Interval: Do not make applications less than 7 days apart.

Maximum Amount per Application: Do not apply more than 11.2 ounces of product or 0.10 pound active ingredient per application.

Maximum Amount of Hero EW Insecticide allowed per Season: Do not apply more than 30 ounces of product or 0.266 pound active ingredient per acre per season.

Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

Remarks

Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons per acre by air and 50 gallons per acre by ground).

One application may be made pre-bloom and a second application may be made post bloom.

For Crown Borer, apply 0.10 pound active per acre post-harvest (fall) or prebloom (spring), as a drench application directed at the crown of plants in a minimum of 200 gallons water / acre. Greater efficacy is observed at higher water gallonages (up to 400 gallons/acre) or in an application prior to a significant rainfall event.

Do not make both pre-bloom foliar and pre-bloom drench applications.

Follow appropriate spray drift precautions on the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

3 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[prebloom](#)

[For Crown Borer: Post-harvest](#)