

## **GRAPE - WEEVIL VINE BLACK, ETC.**

### General Information

#### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zetacypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zetacypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

#### Limitations, Restrictions, and Exceptions

##### GRAPE

##### RESTRICTIONS

Application Interval: Do not make applications less than 7 days apart.

Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

##### Remarks

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 25 gallons by ground and 10 gallons by air).

Follow appropriate spray drift precautions on the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)