

## **FORESTRY USES - BIENNIAL, PERENNIAL BROADLEAF WEEDS AND WOODY PLANTS**

General Information

### **PRODUCT INFORMATION**

Performance of this product may be affected by local conditions, crop varieties, and application method. User should consult local Extension Service, Agricultural Experiment or University Weed Specialists, and state regulatory agencies for recommendations in his area.

Best results are obtained when product is applied to young succulent weeds that are actively growing. The lower specified rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states where control is difficult, use the higher specified rate.

When product is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications if permitted by this label.

Apply 1 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the listed amount of 2,4-D per acre.

For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended. To do so may reduce herbicide's selectivity and could result in crop damage.

Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. This product contains dimethylamine salt of 2,4-D, one of the least volatile forms of 2,4-D.

Do not allow product to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth. Do not use in greenhouses.

NOTE: Herbicide treatment of public water requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit in your state.

Spray equipment used to apply 2,4-D may not be used for any other purpose until thoroughly cleaned by a suitable chemical cleaner.

Spray Preparation: Add the specified amount of product to approximately 1/2 the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue agitation during application until spray tank is empty.

Use in Liquid Nitrogen Fertilizer: Product may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions:

Fill the spray tank approximately 1/2 full with the liquid nitrogen fertilizer. In a separate clean container, mix the amount of product to be used with an equal amount of water. Add the product mixture to the spray tank while agitating. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. **DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER.** Spray mixture must be used immediately and may not be stored.

Note: Pre-mixing the product with an equal amount of water is important.

#### Spot Treatments

To prevent misapplication, apply spot treatments with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 sq ft as indicated in the label.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of 2,4-D AMINE 4. Apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an

area of 1,000 sq ft. Mix the amount of 2,4-D AMINE 4 (fl oz or ml) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of 2,4-D AMINE 4 required for larger areas, multiply the table value (fl oz or ml) by the thousands of sq ft to be treated. An area of 1000 sq ft is approximately 10.5 x 10.5 yards (strides) in size.

Band Application: 2,4-D AMINE 4 may be applied as a band treatment.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

## Limitations, Restrictions, and Exceptions

### FORESTRY USES

Forest site preparation, forest roadsides, brush control, established conifer release (including Christmas trees and reforestation areas)

### SPECIFIC USE DIRECTIONS

Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult-to-control perennial broadleaf weeds and woody species, use up to 8 pts. of 2,4-D AMINE 4 and 1 to 4 qts. of Triclopyr 3A herbicide per acre.

For conifer release, make application in early spring before budbreak of conifers when weeds are small and actively growing.

### FORESTRY PRECAUTIONS:

- Do not allow sprays to contact conifer shoot growth (current year's new growth) or injury may occur.
- For conifer release, do not use on plantations where pine or larch are among the desired species.

### FORESTRY RESTRICTIONS:

- Do not apply to nursery seedbeds.
- Grazing and Haying Restrictions: If grazing or haying is anticipated, do not apply more than 4 pt/acre of 2,4-D AMINE 4 per application. Do not harvest forage or hay from treated areas for 7 days after application.
- Do not make more than one broadcast application per year.
- For broadcast applications, do not apply more than 8 pt/acre of 2,4-D AMINE 4 per 12-month period.

- Basal spray, Cut Surface – Stumps, and Frill:
- Limit of one basal spray or cut surface application per year.
- Maximum of 8.0 lbs ae per 100 gallons of spray solution.
- Injection:
- Limit to one injection application per year.
- Maximum of 2.0 ml of 4.0 lbs ae formulation per injection site.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

48 hours

#### Timings

[When weeds are small and growing actively before the bud stage.](#)