

FIELD ROW CROPS AND VEGETABLES

General Information

General Information

Foli-Zyme is formulated as a soluble concentrate that requires a minimum 1 to 10 dilution with water for all applications.

Foli-Zyme is a fluid nutrient compound designed to prevent nutrient deficiencies in plants.

Foli-Zyme is a valuable supplement to soil- applied nutrients and is particularly beneficial under conditions where soil nutrients are not readily available.

Foli-Zyme will provide nutrients at periods of fruit or flower development.

Foli-Zyme has a pH of 2.0 to 3.5 and will buffer spray tank water. The use of additional buffering agents may result in crop damage.

Foli-Zyme will disperse in water with little agitation. Many pesticides can be added and applied while spraying Foli- Zyme.

Follow this mixing procedure: 1. Water, 2. Foli- Zyme , 3. Pesticide CAUTION: Be sure to conduct "Jar Test" using all products in proper proportion in order to establish physical compatibility.

Recommended for use: Foli- Zyme may be used on the following crops: Alfalfa, Almonds, Apples, Artichokes, Asparagus, Avocados, Barley, All beans, Broccoli, Cabbage, Carrots, Cauliflower, Celery, Citrus, Corn, Cotton, Grapes, Hops, Lettuce, Melons, Milo, Nectarines, Peaches, Peanuts, Peas, Pecans, Peppers, Pistachios, Potatoes, Rice, Soybean, Sorghum, Strawberries, Sugar beets, Sugar cane, Sunflowers, Sweet corn, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Limitations, Restrictions, and Exceptions

Field row crops and vegetables: To supplement soil- applied fertilizer, foliar spray Foli- Zyme on the following schedule: Apply 1 gallon per acre (9 liters per hectare). Use a row- directed spray. Start applications when the crop has 2 to 3 leaves or is

20 days old. Continue sprays at 10 to 14 day intervals.

Method

[Foliar spray](#)

[Directed](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[When the crop has 2 to 3 leaves or is 20 days old.](#)