

TREE, FRUITS AND NUTS

General Information

General Information

Foli-Zyme is formulated as a soluble concentrate that requires a minimum 1 to 10 dilution with water for all applications.

Foli-Zyme is a fluid nutrient compound designed to prevent nutrient deficiencies in plants.

Foli-Zyme is a valuable supplement to soil- applied nutrients and is particularly beneficial under conditions where soil nutrients are not readily available.

Foli-Zyme will provide nutrients at periods of fruit or flower development.

Foli-Zyme has a pH of 2.0 to 3.5 and will buffer spray tank water. The use of additional buffering agents may result in crop damage.

Foli-Zyme will disperse in water with little agitation. Many pesticides can be added and applied while spraying Foli- Zyme.

Follow this mixing procedure: 1. Water, 2. Foli- Zyme , 3. Pesticide CAUTION: Be sure to conduct "Jar Test" using all products in proper proportion in order to establish physical compatibility.

Recommended for use: Foli- Zyme may be used on the following crops: Alfalfa, Almonds, Apples, Artichokes, Asparagus, Avocados, Barley, All beans, Broccoli, Cabbage, Carrots, Cauliflower, Celery, Citrus, Corn, Cotton, Grapes, Hops, Lettuce, Melons, Milo, Nectarines, Peaches, Peanuts, Peas, Pecans, Peppers, Pistachios, Potatoes, Rice, Soybean, Sorghum, Strawberries, Sugar beets, Sugar cane, Sunflowers, Sweet corn, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Limitations, Restrictions, and Exceptions

Tree, fruits and nuts: Apply 1 gallon per acre (9 liters per hectare) 5 to 6 times annually, beginning with delayed dormant spray. Spray intervals of 15 days are suggested for best results. AVOID PERIODS WHEN FRUIT IS EXTREMELY SENSITIVE

TO SPRAY DAMAGE.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Timings

[5 to 6 times annually, beginning with delayed dormant spray.](#)