

POTATOES

General Information

PRODUCT INFORMATION

Revus Top is a broad spectrum product containing two fungicides. It has preventative, systemic and curative properties and is recommended for the control of many important plant diseases. Revus Top provides excellent disease control of many leaf spots, powdery mildews, and downy mildews. Revus Top is applied as a foliar spray and can be used in block, alternating spray, or tank mix programs with other crop protection products. All applications must be made according to the use directions that follow.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Mix only the amount of spray solution needed for immediate application. Avoid spray overlap, as crop injury may occur.

Adjuvants: A spreading/penetrating type adjuvant such as a non-ionic surfactant, crop oil concentrate, or blend is recommended at the manufacturer's recommended rates. When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Revus Top has been used. If fungal isolates that are resistant to Group 3 or Group 40 fungicides are present, efficacy may be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

Integrated Pest Management (IPM): Revus Top should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established

for your area. Revus Top may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

Resistance Management

Revus Top contains two fungicides - mandipropamid, a Carboxylic Acid Amide (CAA) fungicide in Group 40 and difenoconazole, a triazole fungicide in Group 3. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in the label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. SYNGENTA encourages responsible resistance management to ensure effective long-term control of the fungal diseases on the label. Revus Top must not be alternated or tank mixed with any fungicide to which resistance has already developed.

As part of a resistance management strategy:

- Apply a maximum of 4 sprays during one crop cycle
- Apply no more than 2 sequential applications unless otherwise stated in the crop section.
- When tank mixing or alternating, use an effective partner - one that provides satisfactory disease control when used alone at the mixture rate.
- Do not use Revus Top for transplant production.

Application Instructions

Revus Top may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Ground Application:

- Apply in a minimum of 10 gal of water per acre, unless specified otherwise.
- Do not apply through any ultra-low volume (ULV) spray system.
- Thorough coverage is necessary to provide good disease control.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- Thorough coverage is necessary to provide good disease control.
- Apply in a minimum of 5 gallons of water per acre unless specified otherwise.
- Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.
- Do not apply directly to humans or animals.
- Do not apply through any ultra-low volume (ULV) spray system.

Limitations, Restrictions, and Exceptions

POTATOES

Remarks:

Begin applications prior to disease development and continue throughout the season on a 7-10 day interval. Make no more than 2 consecutive applications before switching to another effective fungicide with a different mode of action. If disease pressure is high, use the shortest interval and highest rate. The addition of a spreading/penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is recommended when applying by ground or air.

Application: For best results, use sufficient water volume to provide thorough coverage. Revus Top may be applied by ground, chemigation, or aerial application. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

- Do not apply more than 0.52 lb ai/A/season of mandipropamid containing products.
- Do not apply more than 0.46 lb ai/A/season of difenoconazole containing products.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Prior to disease development.](#)