

CITRUS - CLEMENTINE MANDARINE - FIELD APPLICATIONS

General Information

APPLICATION INSTRUCTIONS

Use only as directed. Read label thoroughly before making applications. N-LARGE contains gibberellic acid which is an extremely potent plant growth regulator. When applying plant growth regulators, deviations from or misuse of the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, result in undesirable effects. Always consult the State Extension Service Specialist in your area for the spray regimen best suited to your conditions. Do not apply to plants under pest, nutritional or water stress. When a range of rates is indicated, use the concentration and spray volume recommended locally by the State Extension Service Specialist. For optimum effectiveness, thoroughly spray entire plant. Spray all parts of the plant or crop to receive desired results. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Discard any unused spray material at the end of each day following local, state or Federal Law. For best results, the water pH should be around neutral and always below 8.5. N-LARGE applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity and no wind) will increase absorption by the plant, thus optimizing effectiveness. Make night- time applications when day- time conditions are not conducive to slow drying conditions. Product persistence: Re- apply N-LARGE if significant rain occurs within 2 hours of application. Compatibility: The N-LARGE spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. Stoller Enterprises does not assume responsibility for unexpected results due to the tank mixing of N-LARGE with other products. DO NOT apply using ULV application methods. For aerial applications, make spray volumes greater than 2 gallons per acre (20 l/ ha), 10 gallons per acre for tree crops (100 l/ha). Apply N-LARGE up to 7 days before harvest.

Limitations, Restrictions, and Exceptions

CITRUS - FIELD APPLICATIONS

Clementine Mandarin (Not for use in California)

To increase fruit set and yield.

APPLICATION TIMING/ INSTRUCTIONS

Make one to two applications from 50% petal fall up to 3 weeks after petal fall. Use a dilute spray with sufficient spray volume for adequate coverage of tree canopy.

Note: The number of applications depends upon amount of desired fruit set. Generally, more fruit will be set by 2 applications, earlier applications, higher rates, and climatic conditions more favorable to set. Differences in the crop strain can also interact with the above factors to affect the degree of fruit set achieved. Reductions in final fruit size can occur as a result of excessive fruit set.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil- incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Make one to two applications from 50% petal fall up to 3 weeks after petal fall.](#)