

FRUIT CROPS - RED TART CHERRY

General Information

APPLICATION INSTRUCTIONS

Use only as directed. Read label thoroughly before making applications. N-LARGE contains gibberellic acid which is an extremely potent plant growth regulator. When applying plant growth regulators, deviations from or misuse of the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, result in undesirable effects. Always consult the State Extension Service Specialist in your area for the spray regimen best suited to your conditions. Do not apply to plants under pest, nutritional or water stress. When a range of rates is indicated, use the concentration and spray volume recommended locally by the State Extension Service Specialist. For optimum effectiveness, thoroughly spray entire plant. Spray all parts of the plant or crop to receive desired results. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Discard any unused spray material at the end of each day following local, state or Federal Law. For best results, the water pH should be around neutral and always below 8.5. N-LARGE applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity and no wind) will increase absorption by the plant, thus optimizing effectiveness. Make night- time applications when day- time conditions are not conducive to slow drying conditions. Product persistence: Re- apply N-LARGE if significant rain occurs within 2 hours of application. Compatibility: The N-LARGE spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. Stoller Enterprises does not assume responsibility for unexpected results due to the tank mixing of N-LARGE with other products. DO NOT apply using ULV application methods. For aerial applications, make spray volumes greater than 2 gallons per acre (20 l/ ha), 10 gallons per acre for tree crops (100 l/ha).Apply N-LARGE up to 7 days before harvest.

Limitations, Restrictions, and Exceptions

RED TART CHERRY (ALL STATES EXCEPT CALIFORNIA)

To maintain and extend high fruiting capacity of tart cherry trees and reduce the

occurrence of “blind” nodes. Treatment will cause bud differentiation, which is apparent the year after application. Therefore, changes in shoot, spur, and flower production will not be evident until two or three years after program initiation. Applications must be applied annually to ensure vegetative development and subsequent yield improvement year after year.

APPLICATION TIMING/ INSTRUCTIONS

Apply one spray 14 to 28 days after bloom. Optimum timing is defined as that stage when 3 to 5 terminal leaves have fully expanded, or, at least 1 to 3 inches of terminal shoot extension has occurred. Use 4 to 18 grams a.i./ acre, depending on tree age and vigor (see label for application rates (Grams a.i./ acre) for Tart Cherry Trees by Age). Apply as a concentrate or dilute spray in sufficient water volume to ensure thorough wetting.

Note: Rates are based on expected normal tree vigor at various ages. Adjust rate according to tree vigor. If trees are vigorous, use lowest recommended rates. Lowest rates must also be used on trees that have been heavily pruned or hedged. Use higher rates for trees low in vigor and weak in shoot and spur production. Excessive application rates will increase vegetative growth at the expense of fruit production the following year. Applications will not improve growth of trees under stress conditions, such as nutritional, moisture, or pest. Best results will be obtained when combined with good cultural practices.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil- incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[14 to 28 days after bloom.](#)