

CELERY

General Information

GENERAL USE INSTRUCTIONS

For best results, X- CYTE should be applied before noon or after 4 p.m. a spreader-sticker (surfactant) cleared for application to growing crops should be used with the product. Store X- CYTE in a cool place and out of direct sunlight. Any spigot or pump put into an X- CYTE drum should be cleaned with disinfectant (bleach and water, etc.) before using, unless the whole drum is to be used at one time. Mix X- CYTE with enough water to get thorough coverage of plant surfaces. X- CYTE is compatible with most other spray materials.

Limitations, Restrictions, and Exceptions

CELERY

1st application: use 2 pints/ acre (2.4 lt/ ha) applied to the seed bed at time of seeding or up to 20 days thereafter.

2nd application: use 2 pints/ acre (2.4 lt/ ha) at the time seedlings are transplanted, see transplanting instructions.

3rd application: use 1 pint/ acre (1.2 lt/ ha) 2-3 weeks after trans planting.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

4 hours

Timings

[1st application: at time of seeding or up to 20 days thereafter.](#)

[2nd application: at the time seedlings are transplanted.](#)

3rd application: 2-3 weeks after transplanting.