

ORNAMENTAL TREES AND HERBACEOUS PLANTS

General Information

GENERAL USE INSTRUCTIONS

For best results, X- CYTE should be applied before noon or after 4 p.m. a spreader-sticker (surfactant) cleared for application to growing crops should be used with the product. Store X- CYTE in a cool place and out of direct sunlight. Any spigot or pump put into an X- CYTE drum should be cleaned with disinfectant (bleach and water, etc.) before using, unless the whole drum is to be used at one time. Mix X- CYTE with enough water to get thorough coverage of plant surfaces. X- CYTE is compatible with most other spray materials.

Limitations, Restrictions, and Exceptions

ORNAMENTAL TREES AND HERBACEOUS PLANTS

Apply 2 pints per acre (2.4 lt/ ha) in transplant water.

Apply 1 pint per acre (1.2 lt/ ha) as a foliar spray when growth begins in the early spring.

Apply 1 pint per acre (1.2 lt/ ha) at bud burst.

Apply 1 pint per acre (1.2 lt/ ha) at bud set.

Apply 1 pint per acre (1.2 lt/ ha) at the end of summer to maintain color through autumn.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

4 hours

Timings

At bud burst.

When growth begins in the early spring.

At bud set.

At the end of summer to maintain color through autumn.