

FIELD CROPS - POTATO - APHID, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Preventive Use

For cutworm, armyworm grape colapsis, or stalk borer control, STEED Insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

Limitations, Restrictions, and Exceptions

FIELD CROPS

POTATOES

Restrictions

- Application Interval: Do not make applications less than 21 days apart.
- Restricted to 2 foliar applications per season.
- Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.
- Leaves cannot be used for food or feed.

Remarks

Apply as required by scouting. Timing and frequency of applications should be

based upon insect populations reaching locally determined economic threshold levels.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 3 gallons by air).

Follow appropriate spray drift precautions on the label.

Armyworm fall; Colorado potato beetle: Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.

For Tuberworm control (adults and larvae) apply prior to harvest or senesce when economic thresholds are met. Only tuberworms that have not bored or tunneled into the plant can be controlled with a foliar application of STEED Insecticide.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Upon insect populations reaching locally determined economic threshold levels.](#)