

VEGETABLE AND SMALL FRUIT CROPS - TOMATO

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Preventive Use

For cutworm, armyworm grape colapsis, or stalk borer control, STEED Insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

Limitations, Restrictions, and Exceptions

VEGETABLE AND SMALL FRUIT CROPS

TOMATO

Restrictions

- Application Interval: Do not make applications less than 10 days apart.
- Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

Remarks

Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.

Thorough coverage is essential to achieve control.

Aphid, Armyworm fall: Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product which is labeled for this pest is recommended for control

Tobacco budworm, Colorado potato beetle: Pyrethroid resistance is common for this pest. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer the resistance management statement in the DIRECTION FOR USE section of the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

1 day

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Upon insect populations reaching locally determined economic thresholds.](#)